

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 November 2000 (30.11.2000)

PCT


(10) International Publication Number
WO 00/71129 A1

- (51) International Patent Classification⁷: **A61K 31/53**, (74) Agents: **BABCOCK, Timothy, J. et al.**; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).
C07D 487/04
- (21) International Application Number: **PCT/US00/13420**
- (22) International Filing Date: **16 May 2000 (16.05.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/135,265 21 May 1999 (21.05.1999) US
60/193,727 31 March 2000 (31.03.2000) US
- (71) Applicant: **BRISTOL-MYERS SQUIBB COMPANY**
[US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— *With international search report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*
- (72) Inventors: **HUNT, John, T.**; 7 Skyfield Drive, Princeton, NJ 08540 (US). **BHIDE, Rajeev, S.**; 156 Barnsbury Road, Langhorne, PA 19047 (US). **BORZILLERI, Robert, M.**; 222 Riverwoods Drive, New Hope, PA 18938 (US). **QIAN, Ligang**; 11 Coventry Lane, Hopewell, NJ 08525 (US).



WO 00/71129 A1

(54) Title: **PYRROLOTRIAZINE INHIBITORS OF KINASES**



(I)

(57) Abstract: The present invention provides compounds of formula (I) and pharmaceutically acceptable salts thereof. The formula (I) compounds inhibit the tyrosine kinase activity of growth factor receptors such as VEGFR-2, FGFR-1, PDGFR, HER-1, HER-2, thereby making them useful as anti-cancer agents. The formula (I) compounds are also useful for the treatment of other diseases associated with signal transduction pathways operating through growth factor receptors.

PYRROLOTRIAZINE INHIBITORS OF KINASES

Related Applications

5 This application claims priority benefit under Title 35 § 119(e) of United States provisional Application No. 60/135,265, filed May 21, 1999, and United States provisional Application No. 60/193,727, filed March 31, 2000.

Field of the Invention

10 This invention relates to compounds that inhibit the tyrosine kinase activity of growth factor receptors such as VEGFR-2, FGFR-1, PDGFR, HER1, and HER2, thereby making them useful as anti-cancer agents. The compounds are also useful in the treatment of diseases, other than cancer, which are associated with signal transduction pathways operating through growth factor receptors such as VEGFR-2.

15

Background of the Invention

Normal angiogenesis plays an important role in a variety of processes including embryonic development, wound healing, obesity and several components of female reproductive function. Undesirable or pathological angiogenesis had been
20 associated with disease states including diabetic retinopathy, psoriasis, cancer, rheumatoid arthritis, atheroma, Kaposi's sarcoma and haemangioma (Fan et al, 1995, Trend Pharmacol. Sci. 16: 57-66; Folkman, 1995, Nature Medicine 1: 27-31). Alteration of vascular permeability is thought to play a role in both normal and pathophysiological processes (Cullinan-Bove et al, 1993,, Endocrinology 133: 829-
25 837; Senger et al, 1993 Cancer and Metastasis Reviews, 12: 303-324).

Receptor tyrosine kinases (RTKs) are important in the transmission of biochemical signals across the plasma membrane of cells. These transmembrane molecules characteristically consist of an extracellular ligand-binding domain connected through a segment in the plasma membrane to an intracellular tyrosine
30 kinase domain. Binding of ligand to the receptor results in stimulation of the receptor-associated tyrosine kinase activity that leads to phosphorylation of tyrosine residues on both the receptor and other intracellular proteins, leading to a variety of cellular responses. To date, at least nineteen distinct RTK subfamilies, defined by

amino acid sequence homology, have been identified. One of these subfamilies is presently comprised by the *fms*-like tyrosine kinase receptor, Flt or Flt1 (VEGFR-1), the kinase insert domain-containing receptor, KDR (also referred to as Flk-1 or VEGFR-2), and another *fms*-like tyrosine kinase receptor, Flt4 (VEGFR-3). Two of these related RTKs, Flt and KDR, have been shown to bind vascular endothelial growth factor (VEGF) with high affinity (De Vries et al, 1992, Science 255: 989-991; Terman et al, 1992, Biochem. Biophys. Res. Comm. 1992, 187: 1579-1586). Binding of VEGF to these receptors expressed in heterologous cells had been associated with changes in the tyrosine phosphorylation status of cellular proteins and calcium fluxes. VEGF, along with acidic and basic fibroblast growth factor (aFGF & bFGF) have been identified as having in vitro endothelial cell growth promoting activity. By virtue of the restricted expression of its receptors, the growth factor activity of VEGF, in contrast to that of the FGFs, is relatively specific towards endothelial cells. Recent evidence indicates that VEGF is an important stimulator of both normal and pathological angiogenesis (Jakeman et al, 1993, Endocrinology, 133: 848-859; Kolch et al, 1995, Breast Cancer Research and Treatment, 36: 139-155) and vascular permeability (Connolly et al, 1989, J. Biol. Chem. 264: 20017-20024).

In adults, endothelial cells have a low proliferation index except in cases of tissue remodeling, such as wound healing and the female reproductive cycle, and adipogenesis. However in pathological states such as cancer, inherited vascular diseases, endometriosis, psoriasis, arthritis, retinopathies and atherosclerosis, endothelial cells are actively proliferating and organizing into vessels. Upon exposure to angiogenic stimuli with growth factors such as VEGF and bFGF, endothelial cells re-enter the cell cycle, proliferate, migrate and organize into a three-dimensional network. The ability of tumors to expand and metastasize is dependent upon the formation of this vascular network.

Binding of VEGF or bFGF to their corresponding receptor results in dimerization, autophosphorylation on tyrosine residues and enzymatic activation. These phosphotyrosine residues serve as "docking" sites for specific downstream signaling molecules and enzymatic activation results in proliferation of endothelial cells. Disruption of these pathways should inhibit endothelial cell proliferation. Disruption of the FGFR-1 pathway should also affect tumor cell proliferation since this kinase is activated in many tumor types in addition to proliferating endothelial


cells. Finally, recent evidence also suggests that disruption of VEGF signaling inhibits endothelial cell migration, a critical process in vascular network formation.

The over-expression and activation of VEGFR-2 and FGFR-1 in tumor-associated vasculature has suggested a role for these molecules in tumor angiogenesis. Angiogenesis and subsequent tumor growth is inhibited by antibodies directed against VEGF ligand and VEGF receptors, and by truncated (lacking a transmembrane sequence and cytoplasmic kinase domain) soluble VEGFR-2 receptors. Dominant mutations introduced into either VEGFR-2 or FGFR-1 which result in a loss of enzymatic activity inhibits tumor growth *in vivo*. Antisense targeting of these receptors or their cognate ligands also inhibits angiogenesis and tumor growth. Recent evidence has elucidated, in part, the temporal requirements of these receptors in tumor growth. It appears that VEGF signaling is critical in early tumor growth and bFGF is more important at a later time associated with tumor expansion.

Other RTKs such as HER1 and HER2 are involved in cell proliferation and are associated with diseases such as psoriasis and cancer. Disruption of signal transduction by inhibition of these kinases would have an antiproliferative and therapeutic effect.

Description of the Invention

In accordance with the present invention, compounds of formula I



their enantiomers, diastereomers, and pharmaceutically acceptable salts, prodrugs and solvates thereof inhibit the tyrosine kinase activity of growth factor receptors such as VEGFR-2. In formula I and throughout the specification, the above symbols are defined as follows:

X and Y are independently selected from O, OCO, S, SO, SO₂, CO, CO₂, NR¹⁰, NR¹¹CO, NR¹²CONR¹³, NR¹⁴CO₂, NR¹⁵SO₂, NR¹⁶SO₂NR¹⁷, SO₂NR¹⁸,

- CONR¹⁹, halogen, nitro, cyano, or X or Y are absent;
- Z is selected from O, S, N, or CR²⁰;
- R¹ is hydrogen, CH₃, OH, OCH₃, SH, SCH₃, OCOR²¹, SOR²², SO₂R²³, SO₂NR²⁴R²⁵,
 CO₂R²⁶, CONR²⁷R²⁸, NH₂, NR²⁹SO₂NR³⁰R³¹, NR³²SO₂R³³, NR³⁴COR³⁵,
 5 NR³⁶CO₂R³⁷, NR³⁸CONR³⁹R⁴⁰, halogen, nitro, or cyano;
- R² and R³ are independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted
 alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heterocyclo,
 substituted heterocyclo, aralkyl, substituted aralkyl, heterocycloalkyl or
 substituted heterocycloalkyl, or when X is halo, nitro or cyano R² is absent or
 10 when Y is halo, nitro or cyano R³ is absent;
- R⁴ and R⁵ are independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl,
 heterocyclo, substituted heterocyclo, aralkyl, substituted aralkyl, or R⁴ and R⁵
 may together form an optionally substituted monocyclic 5-7 membered
 saturated or unsaturated carbocyclic or heterocyclic ring, or an optionally
 15 substituted bicyclic 7-11 membered saturated or unsaturated carbocyclic or
 heterocyclic ring, except that when Z is O or S, R⁵ is absent, or when Z is
 nitrogen, R⁴ and R⁵ are not both hydrogen;
- R⁶ is H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclo, substituted
 heterocyclo, NR⁷R⁸, OR⁹ or halogen;
- 20 R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²¹, R²⁴, R²⁵, R²⁶, R²⁷,
 R²⁸, R²⁹, R³⁰, R³¹, R³², R³⁴, R³⁵, R³⁶, R³⁸, R³⁹ and R⁴⁰ are independently
 selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl,
 substituted aryl, heterocyclo, or substituted heterocyclo;
- R²², R²³, R³³ and R³⁷ are independently selected from the group consisting of alkyl,
 25 substituted alkyl, aryl, substituted aryl, heterocyclo, or substituted
 heterocyclo; and
- R²⁰ is selected from the group consisting of hydrogen, lower alkyl or substituted
 alkyl, or R²⁰ is absent if the carbon to which it is attached is part of an
 unsaturated aryl or heteroaryl ring;
- 30 with the provisos that:
- R² may not be hydrogen if X is SO, SO₂, NR¹³CO₂, or NR¹⁴SO₂
 - R³ may not be hydrogen if Y is SO, SO₂, NR¹³CO₂, or NR¹⁴SO₂.

Listed below are definitions of various terms used to describe this invention. These definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either individually or as part of a larger group.

- 5 The term "alkyl" refers to straight or branched chain unsubstituted hydrocarbon groups of 1 to 20 carbon atoms, preferably 1 to 7 carbon atoms. The expression "lower alkyl" refers to unsubstituted alkyl groups of 1 to 4 carbon atoms.
- The term "substituted alkyl" refers to an alkyl group substituted by, for example, one to four substituents, such as, halo, hydroxy, alkoxy, oxo, alkanoyl, aryloxy, alkanoyloxy, amino, alkylamino, arylamino, aralkylamino, disubstituted amines in which the 2 amino substituents are selected from alkyl, aryl or aralkyl; alkanoylamino, aroylamino, aralkanoylamino, substituted alkanoylamino, substituted arylamino, substituted aralkanoylamino, thiol, alkylthio, arylthio, aralkylthio, alkylthiono, arylthiono, aralkylthiono, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, sulfonamido, e.g. SO_2NH_2 , substituted sulfonamido, nitro, cyano, carboxy, carbamyl, e.g. CONH_2 , substituted carbamyl e.g. CONHalkyl , CONHaryl , CONHaralkyl or cases where there are two substituents on the nitrogen selected from alkyl, aryl or aralkyl; alkoxycarbonyl, aryl, substituted aryl, guanidino and heterocyclos, such as, indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl and the like. Where noted above where the substituent is further substituted it will be with alkyl, alkoxy, aryl or aralkyl.
- 10 The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.
- The term "aryl" refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms in the ring portion, such as phenyl, naphthyl, biphenyl and diphenyl groups, each of which may be substituted.
- 15 The term "aralkyl" refers to an aryl group bonded directly through an alkyl group, such as benzyl.
- The term "substituted aryl" refers to an aryl group substituted by, for example, one to four substituents such as alkyl, substituted alkyl, halo, trifluoromethoxy, trifluoromethyl, hydroxy, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, aralkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, ureido, nitro, cyano, carboxy, carboxyalkyl, carbamyl, alkoxycarbonyl, alkylthiono, arylthiono,
- 20

arylsulfonylamine, sulfonic acid, alkylsulfonyl, sulfonamido, aryloxy and the like.

The substituent may be further substituted by hydroxy, alkyl, alkoxy, aryl, substituted aryl, substituted alkyl or aralkyl.

5 The term "heteroaryl" refers to an optionally substituted, aromatic group for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one heteroatom and at least one carbon atom-containing ring, for example, pyridine, tetrazole, indazole.

The term "alkenyl" refers to straight or branched chain hydrocarbon groups of 2 to 20 carbon atoms, preferably 2 to 15 carbon atoms, and most preferably 2 to 8
10 carbon atoms, having one to four double bonds.

The term "substituted alkenyl" refers to an alkenyl group substituted by, for example, one to two substituents, such as, halo, hydroxy, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, sulfonamido, nitro, cyano, carboxy, carbamyl, substituted
15 carbamyl, guanidino, indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl and the like.

The term "alkynyl" refers to straight or branched chain hydrocarbon groups of 2 to 20 carbon atoms, preferably 2 to 15 carbon atoms, and most preferably 2 to 8 carbon atoms, having one to four triple bonds.

20 The term "substituted alkynyl" refers to an alkynyl group substituted by, for example, a substituent, such as, halo, hydroxy, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, sulfonamido, nitro, cyano, carboxy, carbamyl, substituted carbamyl, guanidino and heterocyclo, e.g. imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl
25 and the like.

The term "cycloalkyl" refers to an optionally substituted, saturated cyclic hydrocarbon ring systems, preferably containing 1 to 3 rings and 3 to 7 carbons per ring which may be further fused with an unsaturated C₃-C₇ carbocyclic ring.

Exemplary groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclododecyl, and adamantyl. Exemplary
30 substituents include one or more alkyl groups as described above, or one or more groups described above as alkyl substituents.

The terms "heterocycle", "heterocyclic" and "heterocyclo" refer to an optionally substituted, fully saturated or unsaturated, aromatic or nonaromatic cyclic group, for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one
5 heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a heteroatom may have 1, 2 or 3 heteroatoms selected from nitrogen atoms, oxygen atoms and sulfur atoms, where the nitrogen and sulfur heteroatoms may also optionally be oxidized and the nitrogen heteroatoms may also optionally be quaternized. The heterocyclic group may be attached at any heteroatom or carbon
10 atom.

Exemplary monocyclic heterocyclic groups include pyrrolidinyl, pyrrolyl, indolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl,
15 piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxazepinyl, azepinyl, 4-piperidonyl, pyridyl, N-oxo-pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1, 1-dioxothienyl, dioxanyl, isothiazolidinyl, thietanyl, thietiranyl, triazinyl, and triazolyl, and the like.

Exemplary bicyclic heterocyclic groups include 2,3-dihydro-2-oxo-1H-indolyl, benzothiazolyl, benzoxazolyl, benzothienyl, quinuclidinyl, quinoliny, quinolinyl-N-oxide, tetrahydroisoquinoliny, isoquinoliny, benzimidazolyl, benzopyranyl, indoliziny, benzofuryl, chromonyl, coumariny, cinnoliny, quinoxaliny, indazolyl, pyrrolopyridyl, furopyridiny (such as furo[2,3-c]pyridiny, furo[3,1-b]pyridiny] or
25 furo[2,3-b]pyridiny), dihydroisoindolyl, dihydroquinazoliny (such as 3,4-dihydro-4-oxo-quinazoliny), benzisothiazolyl, benzisoxazolyl, benzodiaziny, benzofurazany, benzothiopyranyl, benzotriazolyl, benzpyrazolyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, dihydrobenzopyranyl, indoliny, isochromany, isoindoliny, naphthyridiny, phthalaziny, piperonyl, puriny, pyridopyridyl, quinazoliny, tetrahydroquinoliny, thienofuryl, thienopyridyl, thienothienyl, and the like.
30

Exemplary substituents include one or more alkyl or aralkyl groups as described above or one or more groups described above as alkyl substituents.

Also included are smaller heterocyclos, such as, epoxides and aziridines.

The term "heteroatoms" shall include oxygen, sulfur and nitrogen.

The compounds of formula I may form salts which are also within the scope of this invention. Pharmaceutically acceptable (i.e. non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolating or purifying the compounds of this invention.

The compounds of formula I may form salts with alkali metals such as sodium, potassium and lithium, with alkaline earth metals such as calcium and magnesium, with organic bases such as dicyclohexylamine, tributylamine, pyridine and amino acids such as arginine, lysine and the like. Such salts can be formed as known to those skilled in the art.

The compounds for formula I may form salts with a variety of organic and inorganic acids. Such salts include those formed with hydrogen chloride, hydrogen bromide, methanesulfonic acid, sulfuric acid, acetic acid, trifluoroacetic acid, oxalic acid, maleic acid, benzenesulfonic acid, toluenesulfonic acid and various others (e.g., nitrates, phosphates, borates, tartrates, citrates, succinates, benzoates, ascorbates, salicylates and the like). Such salts can be formed as known to those skilled in the art.

In addition, zwitterions ("inner salts") may be formed.

All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The definition of compounds according to the invention embraces all the possible stereoisomers and their mixtures. It very particularly embraces the racemic forms and the isolated optical isomers having the specified activity. The racemic forms can be resolved by physical methods, such as, for example, fractional crystallization, separation or crystallization of diastereomeric derivatives or separation by chiral column chromatography. The individual optical isomers can be obtained from the racemates from the conventional methods, such as, for example, salt formation with an optically active acid followed by crystallization.

Compounds of the formula I may also have prodrug forms. Any compound that will be converted in vivo to provide the bioactive agent (i.e., the compound for formulas I) is a prodrug within the scope and spirit of the invention.

Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, et al. (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);

It should further be understood that solvates (e.g., hydrates) of the compounds of formula I are also within the scope of the present invention. Methods of solvation are generally known in the art.

15 Use and Utility

The present invention is based on the discovery that certain pyrrolotriazines are inhibitors of protein kinases. More specifically, they inhibit the effects of VEGF, a property of value in the treatment of disease states associated with angiogenesis and/or increased vascular permeability such as cancer. The invention relates to a pharmaceutical composition of compound of formula I, or pharmaceutically acceptable salt or hydrate thereof, and a pharmaceutically acceptable carrier in the treatment of hyperproliferative disorder in mammal. In particular, the said pharmaceutical composition is expected to inhibit the growth of those primary and recurrent solid tumors which are associated with VEGF, especially those tumors which are significantly dependent on VEGF for their growth and spread, including for example, cancers of the bladder, squamous cell, head, colorectal, oesophageal, gynecological (such as ovarian), pancreas, breast, prostate, lung, vulva, skin, brain, genitourinary tract, lymphatic system (such as thyroid), stomach, larynx and lung. In another embodiment, the compounds of the present invention are also useful in the treatment of noncancerous disorders such as diabetes, diabetic retinopathy, psoriasis, rheumatoid arthritis, obesity, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies (including proliferative glomerulonephritis and diabetes-induced renal disease), atheroma, arterial restenosis, autoimmune diseases, acute inflammation and

ocular diseases with retinal vessel proliferation, diabetic retinopathy, retinopathy of prematurity and macular degeneration. The invention also relates to prevention of blastocyte implantation in a mammal, treatment of atherosclerosis, excema, sclerodema, hemangioma. Compounds of the present invention posses good activity
5 against VEGF receptor tyrosine kinase while possessing some activity against other tyrosine kinases.

Thus according to a further aspect of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or
10 vascular permeability reducing effect in a warm-blooded animal such as a human being.

According to a further feature of the invention there is provided a method for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal, such as a human being, in need of such treatment which comprises
15 administering to said animal an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof as defined herein before.

The compounds described herein also inhibit other receptor tyrosine kinases including HER1 and HER2 and are therefore useful in the treatment of proliferative disorders such as psoriasis and cancer. The HER1 receptor kinase has been shown to
20 be expressed and activated in many solid tumors including non-small cell lung, colorectal, and breast cancer. Similarly, the HER2 receptor kinase has been shown to be overexpressed in breast, ovarian, lung and gastric cancer. Monoclonal antibodies that downregulate the abundance of the HER2 receptor or inhibit signaling by the HER1 receptor have shown anti-tumor efficacy in preclinical and clinical studies. It
25 is therefore expected that inhibitors of the HER1 and HER2 kinases will have efficacy in the treatment of tumors that depend on signaling from either of the two receptors. The ability of these compounds to inhibit HER1 further adds to their use as anti-angiogenic agents. See the following documents and references cited therein:
Cobleigh, M. A., Vogel, C. L., Tripathy, D., Robert, N. J., Scholl, S., Fehrenbacher,
30 L., Wolter, J. M., Paton, V., Shak, S., Lieberman, G., and Slamon, D. J.,
"Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease", *J. of Clin. Oncol.* 17(9), p.

2639-2648 (1999); Baselga, J., Pfister, D., Cooper, M. R., Cohen, R., Burtness, B., Bos, M., D'Andrea, G., Seidman, A., Norton, L., Gunnett, K., Falcey, J., Anderson, V., Waksal, H., and Mendelsohn, J., "Phase I studies of anti-epidermal growth factor receptor chimeric antibody C225 alone and in combination with cisplatin", *J. Clin.*

5 *Oncol.* 18(4), p. 904-914 (2000).

In addition, the formula I compounds of this invention may be used as contraceptives in mammals.

The antiproliferative, antiangiogenic and/or vascular permeability reducing treatment defined herein before may be applied as a sole therapy or may involve, in
10 addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. The compounds of this invention may also be useful in combination with known anti-cancer and cytotoxic agents and treatments, including radiation. If formulated as a
15 fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. Compounds of formula I may be used sequentially with known anticancer or cytotoxic agents and treatment, including radiation when a combination formulation is inappropriate.

20 In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the antiproliferative, antiangiogenic and/or vascular permeability reducing treatment defined herein before may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three
25 main categories of therapeutic agent:

- (i) antiangiogenic agents that work by different mechanisms from those defined hereinbefore (for example, linomide, inhibitors of integrin $\alpha v \beta 3$ function, angiostatin, razoxin);
- (ii) cytostatic agents such as antiestrogens (for example tamoxifen,
30 toremifen, raloxifene, droloxifene, iodoxifene), progestogens (for example megestrol acetate), aromatase inhibitors (for example anastrozole, letrozole, borazole, exemestane), antihormones, antiprogestogens, antiandrogens (for example flutamide, nilutamide,

bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example gosereline acetate, luprolide), inhibitors of testosterone 5 α -dihydroreductase (for example finasteride), farnesyltransferase inhibitors, anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example EGF, FGF, platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

(iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside); Intercalating antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide nitrosoureas, thiotepan); antimetotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere and newer microtubule agents such as epothilone analogs, discodermolide analogs, and eleutherobin analogs); topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan); cell cycle inhibitors (for example flavopyridols); and biological response modifiers.

As stated above, the formula I compounds of the present invention are of interest for their antiangiogenic and/or vascular permeability reducing effects. Such compounds of the invention are expected to be useful in a wide range of disease states including cancer, diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, obesity, acute and chronic nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation and ocular diseases associated with retinal vessel proliferation such as diabetic retinopathy.

More specifically, the compounds of formula I are useful in the treatment of a variety of cancers, including (but not limited to) the following:

- 5 -carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma;
- 10 -hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burkett's lymphoma;
- hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia;
- 15 -tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma;
- tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas; and
- other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratocanthoma, thyroid
- 20 follicular cancer and Kaposi's sarcoma.

Due to the key role of kinases in the regulation of cellular proliferation in general, inhibitors could act as reversible cytostatic agents which may be useful in the treatment of any disease process which features abnormal cellular proliferation, e.g., benign prostate hyperplasia, familial adenomatosis polyposis, neuro-fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis following angioplasty or vascular surgery, hypertrophic scar formation, inflammatory bowel disease, transplantation rejection, endotoxic shock, and fungal infections.

Compounds of formula I may induce or inhibit apoptosis. The apoptotic response is aberrant in a variety of human diseases. Compounds of formula I, as modulators of apoptosis, will be useful in the treatment of cancer (including but not limited to those types mentioned hereinabove), viral infections (including but not limited to herpevirus, poxvirus, Epstein-Barr virus, Sindbis virus and adenovirus), prevention of AIDS development in HIV-infected individuals, autoimmune diseases

(including but not limited to systemic lupus, erythematosus, autoimmune mediated glomerulonephritis, rheumatoid arthritis, psoriasis, inflammatory bowel disease, and autoimmune diabetes mellitus), neurodegenerative disorders (including but not limited to Alzheimer's disease, AIDS-related dementia, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, spinal muscular atrophy and cerebellar degeneration), myelodysplastic syndromes, aplastic anemia, ischemic injury associated with myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, hematological diseases (including but not limited to chronic anemia and aplastic anemia), degenerative diseases of the musculoskeletal system (including but not limited to osteoporosis and arthritis) aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases and cancer pain.

The compounds of formula I are especially useful in treatment of tumors having a high incidence of tyrosine kinase activity, such as colon, lung, and pancreatic tumors. By the administration of a composition (or a combination) of the compounds of this invention, development of tumors in a mammalian host is reduced.

Compounds of formula I may also be useful in the treatment of diseases other than cancer that may be associated with signal transduction pathways operating through growth factor receptors such as VEGFR-2.

The compounds of this invention may be formulated with a pharmaceutical vehicle or diluent for oral, intravenous or subcutaneous administration. The pharmaceutical composition can be formulated in a classical manner using solid or liquid vehicles, diluents and additives appropriate to the desired mode of administration. Orally, the compounds can be administered in the form of tablets, capsules, granules, powders and the like. The compounds may be administered in a dosage range of about 0.05 to 200 mg/kg/day, preferably less than 100 mg/kg/day, in a single dose or in 2 to 4 divided doses.

Biological assays

VEGFR-2 and FGFR-1 Kinase assays:

| Reagents | Final Concentration | |
|-----------------------|---------------------|---------------|
| | <u>VEGFR-2</u> | <u>FGFR-1</u> |
| <u>Stock Solution</u> | | |
| Tris pH 7.0 | 20 mM | 20mM |
| BSA 10 mg/ml | 25 µg/ml | 25µg/ml |

| | | | |
|---|---------------------------------------|------------|----------|
| | MnCl ₂ (1M) | 1.5 mM | 0.5 mM |
| | MgCl ₂ (1M) | ----- | 0.5 mM |
| | DTT(1M) | 0.5 mM | 0.5 mM |
| | Enzyme Stock in 10%glycerol (1 mg/ml) | 5ng/rxn | 20ng/rxn |
| 5 | Polyglu/tyr (10 mg/ml) | 80 µg/ml | 30 µg/ml |
| | ATP (1mM) | 2.5 µM | 1.0 uM |
| | γ-ATP (10µCi/µl) | 0.5 µCi/ml | 0.5µCi |

Incubation mixtures employed for VEGFR-2 or FGFR-1 assay contain the synthetic substrate polyGlu:Tyr, (4:1), ATP, ATP-γ-³³P and buffer containing Mn⁺⁺ and/or Mg⁺⁺, DTT, BSA, and Tris buffer. The reaction is initiated by addition of enzyme and after 60 minutes is terminated by the addition of TCA to 30%. Inhibitors are brought to 10mM in 100% DMSO. Assays are prepared in a 96 well format. Compounds are diluted 1:500 in 100% DMSO and then 1:10 in water for a final DMSO concentration of 10%. 10 µL are added to rows B-H in a 96 well format of 10% DMSO. 20 µl of compound is added to row A at a concentration 5 fold higher than running conditions. Ten µL are transferred to each row with 10 pipetting phases for mixing, and at row F 10 µL are discarded. Row G is a control with no compound and row H is no compound and no enzyme control. Enzyme and substrate are delivered using a Tomtec Quadra station.

Plates are covered with sticky plate tops, incubated at 27°C for 60 minutes, and then acid precipitated with TCA for 20 minutes on ice. The precipitate is transferred to UniFilter-96, GF/C microplates using either a Tomtec or Packard FilterMate harvester. Activity is determined by quantitating the incorporated radioactivity using a Packard TopCount Microplate Scintillation Counter following the addition of Microscint-20 cocktail into each dried well of the UniFilter microplates.

The instant compounds inhibit VEGFR-2 and FGFR-1 kinases with IC₅₀ values between 0.003 -25 µM.

HER1 or HER2 Kinase assays:

Compounds of interest were assayed in a kinase buffer that contained 20 mM Tris.HCl, pH 7.5, 10 mM MnCl₂, 0.5 mM dithiothreitol, bovine serum albumin at 0.1 mg/ml, poly(glu/tyr, 4:1) at 0.1 mg/ml, 1 µM ATP, and 4 µCi/ml [γ-³³P]ATP.

Poly(glu/tyr, 4:1) is a synthetic polymer that serves as a phosphoryl acceptor and is purchased from Sigma Chemicals. The kinase reaction is initiated by the addition of enzyme and the reaction mixtures were incubated at 26 °C for 1 h. The reaction is terminated by the addition of EDTA to 50 mM and proteins are precipitated by the addition of trichloroacetic acid to 5%. The precipitated proteins are recovered by filtration onto Packard Unifilter plates and the amount of radioactivity incorporated is measured in a Topcount scintillation counter.


For the preparation of recombinant HER1, the cytoplasmic sequence of the receptor were expressed in insect cells as a GST fusion protein, which was purified by affinity chromatography. The cytoplasmic sequence of HER2 was subcloned into the baculovirus expression vector pBlueBac4 (Invitrogen) and was expressed as an untagged protein in insect cells. The recombinant protein was partially purified by ion-exchange chromatography.

The instant compounds inhibit HER-1 and HER-2 kinases with IC₅₀ values between 0.003 -25 μM.

Methods of Preparation

Certain compounds of formula I may generally be prepared according to the following schemes and the knowledge of one skilled in the art.

Scheme 1

**Step 1**

The first step is accomplished by the reaction of an optionally substituted 2-formylpyrrole (product 1) with an aminating reagent, such as hydroxylamine-O-sulfonic acid, in an aqueous solvent at room temperature, followed by treatment under cooling with a base such as KOH . Compounds 1 may be obtained from substituted pyrroles by formylation, for example by reaction with phosphorus oxychloride and DMF. A methylpyrrole may be obtained by reduction of a formylpyrrole, for example by reaction with lithium aluminum hydride.

Step 2

The product 2 is reacted with an aqueous base such as KOH at room temperature to form the product 3 of Scheme 1.

Step 3

The compound 3 is reacted with an acylating agent, such as formic acid, in an aqueous solvent, to form the product 4 of Scheme 1.

Step 4

The compound 4 is cyclized with a base such as sodium methoxide in methanol with heating to form the product 5 of Scheme 1.

Step 5


The compound 5 is halogenated, for example with phosphorus oxybromide at elevated temperature, to form the product 6 of Scheme 1.

Step 6

- 5 The compound 6 is reacted with an amine such as an aniline in an organic solvent, such as acetonitrile, to form the product 7 of Scheme 1.

The compound 7 of Scheme 1 where R_1 = 7-halogen can be prepared from the compound 7 of Scheme 1 where R_1 = hydrogen by reaction with a halogenating agent such as bromine in a suitable solvent such as acetic acid.

10

Scheme 2

E = electron withdrawing group such as ester or nitro or ketone

X^I = halogen

Step 1

- 15 An anion of tosylmethyl isocyanide (TosMIC) is reacted with a Michael acceptor such as ethyl crotonate to obtain disubstituted pyrrole 1. An anion of TosMIC could be made by treating a solution of it in dimethyl sulfoxide (DMSO) with a base such as sodium hydride (NaH) at rt or a solution of it in tetrahydrofuran (THF) with lithium hexamethyldisilazane at -78°C .

Step 2

Treatment of pyrrole 1 with an acylating agent such as trichloroacetyl chloride in the presence of a Lewis acid such as aluminum chloride at from rt to 50 °C followed by treatment with sodium methoxide could afford trisubstituted pyrrole 2. Alternatively, following the published procedure of (M. Suzuki, M. Miyoshi, K. Matsumoto J. Org. Chem. **1974**, 39, 1980) Compound 2 could be obtained by warming an aldehyde, such as acetaldehyde, with 2 equivalents of ethyl isocyanoacetate in the presence of a base, such as DBU, in an organic solvent, such as THF.

Step 3

10 Pyrrole 2 could be aminated by an aminating reagent, such as diphenyl phosphoryl hydroxylamine, in the presence of a base, such as sodium hydride at rt in organic solvents, such as dimethyl formamide (DMF).

Step 4

15 N-Aminated pyrrole 3 upon heating at from 120 to 195 °C with formamide could undergo cyclization to afford 1,2,4-triazine 4.

Step 5


Compound 4 upon treatment with a halogenating agent, such as phosphorous oxybromide at from 60 to 115 °C, in the presence or absence of a co-solvent such as 1,2-dichloroethane, compound 5 could be obtained.

20 Step 6

Compound 5 is reacted with amines, such as anilines in an organic solvent, such as DMF, to obtain compound 6. Alternatively, compound 5 is treated with an anion of a heterocyclic compound, such as oxindole, in an organic solvent such as THF.

25

Scheme 3



wherein $R^a = XR^2$, $R^b = R^6$ described hereinbefore

Step 1

A suitably N-protected ester of glycine, such as with benzyl group, could be added to dialkyl methylene malonate at from rt to 80 °C to obtain compound 1.

5 Step 2

Compound 1 could undergo cyclization to form pyrrole 2 upon treatment with a strong base, such as lithium hexamethyldisilazane at from -78 °C to rt in an organic solvent such as THF.

Step 3

10 Compound 2 could be alkylated by treatment with an alkylating agent, such as iodomethane or dimethyl sulfate, in the presence of a base, such as potassium carbonate, in an organic solvent, such as acetone or DMF.

Step 4

15 Deprotection of compound 3 could be achieved, when optionally protected by groups such as benzyl, by hydrogenation over a catalyst, such as palladium, in the presence of ammonium formate.


Step 5

Compound 4 could be converted to compound 5 in an analogous manner to that described in Scheme 2.

20 Step 6

Hydrolysis of the ester group in compound 5 could be achieved by treatment with a base such as aqueous potassium hydroxide. The resulting acid could be coupled with an amine in the presence of a coupling agent, such as DCC or PyBrop.

Scheme 4



wherein $X = NR^{10}$, $NR^{11}CO$, $NR^{12}CONR^{13}$, $NR^{14}COO$, $NR^{15}SO_2$, $NR^{16}SO_2NR^{17}$, as described hereinbefore.

Step 1

Compound 5 of Scheme 3 could be converted to carboxylic acid by treatment with a base such as aqueous potassium hydroxide. This acid could undergo Curtius rearrangement by treatment with diphenyl phosphoryl azide in the presence of an alcohol, such as benzyl alcohol, in an organic solvent, such as 1,4-dioxane, to afford compound 1.


Step 2

Deprotection of the carbamate group could be achieved, when optionally protected by groups such as carbobenzyloxy, by hydrogenation over a catalyst, such as palladium.

Step 3

The amino group of compound 2 could be acylated, for example by treatment with a carboxylic acid in the presence of a coupling agent such as DCC, or could be sulfonylated, for example by treatment with a sulfonyl chloride. Alternatively, the amino group of compound 2 may be alkylated with alkyl halides or could undergo reductive amination with aldehydes in the presence of a reducing agent, such as sodium cyanoborohydride.

Scheme 5



wherein $R^a = XR^2$; $R^c, R^d = R^6$; and $R^1 = H$ or $COOR^{26}$ as described hereinbefore.

Step 1

- 5 Suitably protected compound 1 (imino dicarboxylate) could be cyclized by treatment with dialkyl oxalate in the presence of a base, such as sodium methoxide, in an organic solvent, such as methanol.

Step 2

- 10 Compound 2 upon selective deprotection, such as with trifluoro acetic acid (TFA) when optionally protected by tert-butyl ester, undergoes decarboxylation to afford Compound 3 where $R^1 = H$. Step 2 is omitted to form compound 3 where $R^1 = COOR^{26}$.

Step 3

- 15 The hydroxy group of compound 3 could be etherified by reaction with an alkylating agent, such as dimethyl sulfate.


Step 4

Compound 4 could be deprotected by hydrogenation, when optionally protected as benzyl group, to obtain compound 5.

Step 5

- 20 Compound 5 could then be converted to compound 6 in an analogous manner to that described in step 4 of scheme 3 followed by steps 3 to 6 of Scheme 2.

Scheme 6

**Step 1**

Compound 5 of Scheme 2 could be etherified at the 4-position, for example by treatment with phenoxide anion.

Step 2

Reduction with a reducing agent, such as DIBAL, in an organic solvent, such as toluene, could afford the alcohol 2.

Step 3

Oxidation of the alcohol could be achieved by treatment of compound 2, for example with MnO₂ at an elevated temperature in an organic solvent, such as toluene.

Step 4

Treatment of compound 3 with an oxidant, such as m-chloroperbenzoic acid (m-CPBA) in an organic solvent, such as dichloromethane, followed by aqueous

hydrolysis with a base, such as potassium bicarbonate, could afford the hydroxy compound 4.

Step 5


Alkylation of the phenolic group in compound 4 with an agent, such as iodomethane, in the presence of a base, such as NaH, at from rt to 100 °C, could afford compound 5.

Step 6

Hydrolysis of Compound 5 could be achieved by treatment with an acid, such as aqueous HCl, at an elevated temperature to afford compound 6.

10 Step 7

Compound 6 could be converted to compound 7 using procedures analogous to those described in Scheme 2.



15 Step 1

A compound 3 of Scheme 6 could undergo Wittig reaction, for example with phosphonates such as methyl diethylphosphonoacetate, in an organic solvent, such as dichlorethane, in the presence of a base, such as NaH to afford Compound 1.

Step 2

The double bond could be hydrogenated by treatment with hydrogen in the presence of a catalyst, such as palladium.

Step 3


5 Compound 2 could be converted to compound 3 by procedures described in Scheme 2.

Step 4

10 Hydrolysis of the ester, as described hereinbefore, followed by coupling of the resulting acid with an amine in the presence of a coupling agent, such as DCC, could afford Compound 4.

In addition, other compounds of formula I may be prepared using procedures generally known to those skilled in the art. In particular, the following examples provide additional methods for the preparation of the compounds of this invention.

15 The invention will now be further described by the following working examples(s), which are preferred embodiments of the invention. All temperatures are in degrees Celsius (°C) unless otherwise indicated. HPLC purifications were done on C18 reverse phase (RP) columns using water methanol mixtures and trifluoroacetic acid as buffer solution. These examples are illustrative rather than limiting and it is to be understood that there may be other embodiments that fall within the spirit and scope of
20 the invention as defined by the claims appended hereto.

Example 1**N-(4-Chlorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine****25 A. 4-Bromo-pyrrolo[2,1-f][1,2,4]triazine**

A mixture of 50 mg (0.37 mmol) of pyrrolo[2,1-f][1,2,4]triazin-4(3H)-one [prepared as described in S.A.Patil, B.A. Otter and R.S. Klein, *J. Het. Chem.*, 31, 781-786 (1994)] and 0.5 g of phosphorus oxybromide was heated at 60°C for 20 min., under argon. A clear orange melt was initially obtained which solidified to a yellow solid


on continued heating. Ice was added with vigorous stirring to the solid. The mixture was extracted twice with ethyl acetate. The combined extracts were washed with sat. NaHCO_3 and brine, dried (MgSO_4), and the solvent removed to afford 63 mg of crude Compound A as an orange oil which crystallized on standing.

5 $(\text{M}+\text{H})^+ = 198^+, 200^+$

B. 4-(4-Chlorophenylamino)-pyrrolo[2,1-f][1,2,4]triazine

A solution of 60 mg (0.3 mmol) of Compound A and 38 mg (0.3 mmol) of p-chloroaniline in 1.5 ml of acetonitrile was stirred overnight at rt and under argon. A white precipitate was obtained which was removed by filtration. The filter cake was
10 suspended in ethyl acetate and sat. NaHCO_3 added and the mixture stirred until a solution was obtained. The organic layer was separated and washed with brine, dried (MgSO_4) and the solvent removed to yield 23 mg (0.094 mmol, 31 %) of Example 1 as a white solid. $(\text{M}+\text{H})^+ = 245^+$

Example 2




15

2-Methyl-5-(pyrrolo[2,1-f][1,2,4]triazin-4-ylamino)phenol

The compound of Example 2 was prepared as a white solid in 49% yield from Compound A of Example 1 and 3-hydroxy-4-methylaniline as described for Compound B of Example 1. $(\text{M}+\text{H})^+ = 241^+$

20


Example 3



N-(4-Chloro-2-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine


The compound of Example 3 was prepared as a white solid in 40% yield from Compound A of Example 1 and 2-fluoro-4-chloroaniline as described for Compound B of Example 1. $(M+H)^+ = 263^+$

5

Example 4**7-Bromo-N-(4-chlorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine**

To a solution of 26 mg (0.11 mmol) of Example 1 in 1 ml of acetic acid was added dropwise a solution of 19 mg of bromine in 100 ml of acetic acid. A white precipitate was obtained during the addition. Stirring was continued for 1 hr, under argon. The mixture was evaporated to dryness and the residue diluted with ethyl acetate and treated with sat NaHCO_3 . The clear colorless organic layer was washed with brine, dried (MgSO_4) and the solvent removed to give a white solid residue. This material was subjected to flash chromatography on a 15 cc column of silica gel. Elution with chloroform (100 %) afforded 18 mg (0.06 mmol, 50%) of Example 4 as a white solid. $(M+H)^+ = 323, 325^+$

20

Example 5**2-Methyl-5-[(6-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol****A. 2-Formyl-3-methylpyrrole/2-formyl-4-methylpyrrole**

To 3.15 ml (41 mmol) of DMF at 0°C under argon was added dropwise 3.81 ml (41 mmol) of phosphorus oxychloride. The cooling bath was removed and stirring was continued for 15 min. The solution was diluted with 9 ml of 1,2-dichloroethane and again cooled to 0°C . A solution of 3.0 g (37 mmol) of 3-methylpyrrole in 9 ml of 1,2-dichloroethane was added dropwise. The mixture was heated to reflux for 15 min,

cooled to 0°C and a solution of 16.2 g (203 mmol) of sodium acetate in 45 ml of water was added with vigorous stirring. The mixture was heated at reflux for 20 min and allowed to cool to room temperature. The aqueous layer was separated and extracted twice with methylene chloride. The combined organic layers were washed with sat
 5 NaHCO₃ until pH 7, dried (MgSO₄), and the solvent removed to yield a dark oily solid which was purified by flash chromatography (10% EtOAc:hexane) to afford 3.6 g (89%) of a 4:1 mixture of 2-formyl-3-methylpyrrole and 2-formyl-4-methylpyrrole as a pale yellow solid.

B. 1-Amino-2-aminocarbonyl-4-methyl-pyrrole

10 Compound A as an isomeric mixture was aminated as described in S.A. Patil, B.A. Otter and R.S. Klein, *J. Het. Chem.*, 31, 781-786 (1994) to form a 2:1 mixture of 1-amino-2-cyano-3-methylpyrrole and 1-amino-2-cyano-4-methylpyrrole in combined 20% yield. The mixture of nitriles was hydrolyzed as described in the reference to form Compound B as well as unreacted 1-amino-2-cyano-3-methylpyrrole, which
 15 were separated by flash chromatography (10% EtOAc:hexane).


C. 6-Methyl-pyrrolo[2,1-f][1,2,4]triazin-4(3H)-one

Compound C was prepared from Compound B as described in S.A. Patil, B.A. Otter and R.S. Klein, *J. Het. Chem.*, 31, 781-786 (1994).

D. 4-(3-Hydroxy-4-methyl-phenylamino)-6-methyl-pyrrolo[2,1-f][1,2,4]triazine

20 Example 5 was prepared from Compound C using the 2 step sequence of Compound A of Example 1 and Compound B of Example 1, using 3-hydroxy-4-methylaniline. (M+H)⁺ = 255.


Example 6



N-(4-Bromo-2-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine

25 The title compound was prepared as a white solid in 57% yield from Compound A of Example 1 and 2-fluoro-4-bromoaniline as described for Compound B of Example 1. (M+H)⁺ = 307, 309.

Example 7



2-Methyl-5-[(5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol

A. 1-Amino-2-aminocarbonyl-3-methyl-pyrrole


A solution of 290 mg (2.4 mmol) of 1-amino-2-cyano-3-methylpyrrole (prepared as described in Compound B of Example 5) and 3.5 g (62 mmol) of potassium hydroxide in 2 ml of water and 28 ml of ethanol was heated at reflux for 3 hr. The mixture was evaporated to near dryness, the residue diluted with additional water, and the mixture extracted with ethyl acetate (3x). The combined extracts were washed with brine, dried (MgSO₄) and the solvent removed to yield 274 mg (82 %) of Compound A as a white solid.

B. 4-(3-Hydroxy-4-methyl-phenylamino)-5-methyl-pyrrolo[2,1-f][1,2,4]triazine

Example 7 was prepared from Compound A as described for Compounds C and D of Example 5. (M+H)⁺ = 255.

15

Example 8




N-(4-Bromo-2-fluorophenyl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine

Example 8 was prepared as a white solid in 29% overall yield from 5-methylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one using the 2 step sequence of Compounds A and B of Example 1, using 2-fluoro-4-bromo-aniline. (M+H)⁺ = 321, 323.

20

Example 9




N-(4-Bromo-2-fluorophenyl)-6-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine

Example 9 was prepared as a white solid in 20% overall yield from Compound C of Example 5 using the 2 step sequence of Compound A and Compound B of Example 1, using 2-fluoro-4-bromo-aniline. The product was purified by flash chromatography on silica gel with 10% ethyl acetate/hexanes. $(M+H)^+ = 321, 323$.

5

Example 10




1-[2,3-Dihydro-6-(pyrrolo[2,1-f][1,2,4]triazin-4-ylamino)-1H-indol-1-yl]ethanone

Example 10 was prepared as a white solid in 4% yield from Compound A of Example 1 and 1-acetyl-6-aminoindoline as described for Compound B of Example 1, using 10% isopropanol/methylene chloride for extraction, and with purification by chromatography on silica gel with ethyl acetate. $(M+H)^+ = 294$.

10

Example 11



2-Methyl-5-[(7-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol

15

A. 2-Methylpyrrole

To 80 ml of a 1 M solution of lithium aluminum hydride in ether (80 mmol), at rt and under argon, was added dropwise a solution of 4.0 g (40 mmol) of 2-formylpyrrole at such a rate as to maintain gentle reflux. Reflux was continued for 6 hr. The excess hydride was hydrolyzed by the dropwise sequential addition of 3 ml of water (with cooling), 3 ml of 15% sodium hydroxide, and 9 ml of water. The resulting solids were removed by filtration and the filter cake washed well with additional ether. The filtrate was evaporated to dryness and the dark oil residue diluted with methylene chloride, dried ($MgSO_4$) and the solvent removed. The resulting dark oil was distilled (kugelrohr, 700 mm, $100^\circ C$) to afford 1.13 g (35%) of Compound A as a clear colorless oil.

25

B. 2-Formyl-5-methyl-pyrrole

To 0.54 ml of DMF, with ice cooling and under argon, was added dropwise 0.64 ml of POCl₃. The reaction, which solidified during the addition, was warmed lightly with a warm water bath and the clear colorless solution stirred an additional 20 min after addition was completed. The mixture was diluted with 3 ml of 1,2-dichloroethane and, with cooling, a solution of 510 mg (6.3 mmol) of Compound A was added dropwise. The solution was heated at reflux for 15 min, ice cooled and a solution of 2.6g (31.5 mmol) of sodium acetate was added with vigorous stirring. The mixture was heated at 80 °C for 20 min, cooled to rt and extracted with methylene chloride. The extracts were washed with brine, dried (MgSO₄) and the solvent removed to give a dark oil residue. This material was subjected to flash chromatography on silica with 10% EtOAc:hexane to afford 126 mg (18%) of Compound B as a light tan solid.

C. 1-Amino-2-cyano-5-methyl-pyrrole

A solution of 140 mg (1.28 mmol) of Compound B in 2 ml of methylene chloride, at rt and under argon, was added rapidly dropwise to a solution of 290 mg (1.35 mmol) of MSH (O-mesitylenesulfonyl-hydroxylamine) in 2 ml of methylene chloride. A deep red solution was obtained which was stirred for an additional 0.5 hr. The mixture was washed twice with sat NaHCO₃, dried (MgSO₄) and the solvent removed to give a deep red oil. A solution of this material in 5 ml of DMF was added dropwise to a suspension of 102 mg (2.56 mmol) of sodium hydride in 5 ml of DMF, at rt and under argon. The red color was consumed and the reaction became dark yellow. Stirring was continued for an additional 0.5 hr. The mixture was ice cooled and a solution of 385 mg (1.8 mmol) of MSH in 5 ml of DMF was added. Stirring was continued for 1 hr with cooling. The dark solution was diluted with ethyl acetate and washed once with water. The aqueous layer was reextracted once with ethyl acetate. The combined organic layers were dried (MgSO₄) and the solvents removed. The resulting dark red oil was subjected to flash chromatography on silica with 20% EtOAc:hexane to afford 100 mg (65%) of Compound C as a yellow solid.

D. 1-Amino-2-aminocarbonyl-5-methyl-pyrrole

To a solution of 1.1g (20 mmol) of KOH in 2.5 ml of water was added 100 mg (0.82 mmol) of Compound C. The suspension was stirred overnight at rt and heated at 50°C for 4 hrs. The resulting precipitate was removed by filtration. The filter cake was dissolved in ethyl acetate, the solution was dried (MgSO₄) and the solvent removed to give 58 mg of Compound D as a yellow solid. The aqueous filtrate was

extracted 3x with ethyl acetate. The combined extracts were dried (MgSO_4) and the solvent removed to give an additional 11 mg of Compound D as a yellow solid (total 69 mg, 60%).

E. 7-methyl-pyrrolo[2,1-f][1,2,4]triazin-4(3H)-one

- 5 A solution of 65 mg (0.47 mmol) of Compound D in 0.5 ml of formic acid was heated at 100°C for 4 hrs. After removal of the solvent, the residue was subjected to flash chromatography on silica with 50% EtOAc:hexane to afford 53 mg (76%) of Compound E as a white solid.


F. 4-(3-Hydroxy-4-methyl-phenylamino)-7-methyl-pyrrolo[2,1-f][1,2,4]triazine

10

Example 11 was prepared as a white solid in 30% overall yield from Compound E using the 2 step sequence of Compound A of Example 1 and Compound B of Example 1, using 3-hydroxy-4-methylaniline. The product was purified by flash chromatography on silica with 25% ethyl acetate/hexanes. $(\text{M}+\text{H})^+$ 255.

15

Example 12




N-(4-Bromo-2-fluorophenyl)-7-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine

- Example 12 was prepared as a white solid in 33% overall yield from Compound E of Example 11 using the 2 step sequence of Compound A of Example 1 and Compound B of Example 1, using 2-fluoro-4-bromoaniline. The product was purified by flash chromatography on silica with 10% ethyl acetate/hexanes. $(\text{M}+\text{H})^+ = 321, 323$.

20

Example 13



25

5-[(5,7-Dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol

A. 2-formyl-3,5-dimethylpyrrole

To dimethylformamide (4.5 mL, 57.8 mmol) under argon at 0°C was added phosphorus oxychloride (57.8 mmol) dropwise over 5 min. The cooling bath was removed and after 15 min. 1,2-dichloroethane (15 mL) was added. The reaction mixture was again cooled to 0°C and a solution of 2,4-dimethylpyrrole (52.6 mmol) in 1,2-dichloroethane (15 mL) was added dropwise over 15 min. The reaction was heated to reflux for 15 min, and then cooled to rt. A solution of sodium acetate (24 g) in water (75 mL) was added slowly to the reaction mixture and the resulting mixture was again heated to reflux for 20 min. After the reaction mixture was cooled to rt it was diluted with CH₂Cl₂, and the aqueous phase was washed with CH₂Cl₂ (2x50 mL). The combined organic fractions were washed with saturated NaHCO₃, dried (Na₂SO₄), and concentrated *in vacuo*. The crude material was purified by chromatography on silica gel eluting with 10% ethyl acetate in hexane to provide 5.2 g (80%) of the desired compound 2-formyl-3,5-dimethylpyrrole. [M+H]⁺=124.1, [M-H]⁻=122.0


B. 5,7-dimethylpyrrolo[2,1-f][1,2,4]triazine-4(3H)-one

Compound B was prepared from A as described in [S.A.Patil, B.A. Otter and R.S. Klein, *J. Het. Chem.*, 31, 781-786 (1994)]. Thus, removal of formic acid after the reaction gave 5 mg (95%) of compound B. ¹H NMR (CD₃OD): δ 7.40 (s, 1H), 2.15 (s, 1H), 2.28 (s, 3H), 2.19 (s, 3H)

C. 5,7-Dimethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine

The title compound was prepared by treating compound B with 3-hydroxy-4-methylaniline as described for compound B of Example 1. Thus, after purification by preparative HPLC, 14 mg (25 %) of 5,7-Dimethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine was obtained as white solid. MS: [M+H]⁺=269.2, ESI [M-H]⁻=267.0; ¹H NMR (CDCl₃): δ 8.20 (br s, 1H), 7.81 (s, 1H), 7.06 (d, *J*=8.0 Hz, 1H), 6.99 (s, 2H), 6.69 (dd, *J*=8.0, 5.7 Hz, 1H), 6.25 (s, 1H), 2.58 (s, 3H), 2.43 (s, 3H), 2.07 (s, 3H)

Example 14




5-[(6-Ethyl-5,7-dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol

The title compound was prepared from 2,4-dimethyl-3-ethylpyrrole in a manner similar to the preparation of Example 13 from 3,5-dimethylpyrrole. Thus, after

5 purification by preparative HPLC, 19 mg (21 %) of 5,7-dimethyl-6-ethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine was obtained as white solid. MS: $[M+H]^+ = 297.3$, $[M-H]^- = 295.1$; 1H NMR ($CDCl_3$): δ 9.67 (br s, 1H), 7.81 (s, 1H), 7.10 (d, $J=8.4$ Hz, 1H); 6.90 (br s, 1H), 6.70 (s, 1H), 6.64 (dd, $J=8.3, 5.8$ Hz, 1H), 2.64 (q, $J=7.5$, 1H), 2.55 (s, 3H), 2.44 (s, 3H), 2.08 (s, 3H), 1.13 (t, $J=7.5$, 3H)

10

Example 15



5-[(5,6-Dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol

A. Preparation of 3,4-dimethylpyrrole


To a flask containing anhydrous diethyl ether (160 mL) was added slowly a solution of $LiAlH_4$ in THF (1 M, 71.01 mmol). This mixture was stirred for 15 min. at rt. 15 diethyl 3,4-pyrroledicarboxylate (5.0 g, 23.67 mmol) was added portion-wise as a solid to the solution over 15 min. The reaction mixture was then heated at 45°C (reflux) overnight. The reaction mixture was cooled to 0°C and $Na_2SO_4 \cdot 10 H_2O$ (15 g) was slowly added. The mixture was stirred for 30 min. and 15 mL of 10% aqueous 20 NH_4Cl was added slowly and the mixture stirred for an additional 15 min. The mixture was filtered and the residue was rinsed with ethyl acetate (3 x 20 mL). The filtrate was dried over Na_2SO_4 and purified by chromatography on silica gel eluting with a gradient of 10-20% ethyl acetate in hexanes to provide 540 mg (24%) of 3,4-dimethylpyrrole. MS: $[M-H]^- = 94.0$

B. 5,6-Dimethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine.

The compound A was converted to the title compound in a manner similar to the preparation of Example 13 from 3,5-dimethylpyrrole. Thus, after purification by preparative HPLC, 19 mg (21 %) of 5,6-dimethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine was obtained as white solid. $[M+H]^+ = 269.2$, $[M-H]^- = 267.0$; $^1\text{H NMR}$ (CDCl_3): δ 8.30 (br s, 1H), 7.54 (s, 1H), 7.14 (s, 1H), 6.87 (d, $J=7.9$ Hz, 1H), 6.85 (s, 1H), 6.45 (dd, $J=7.8, 1.9$ Hz, 1H), 2.28 (s, 3H), 1.97 (s, 3H), 1.85 (s, 3H)

10

Example 16



5-[(5-Ethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol

A. Preparation of 2-formyl-3-ethylpyrrole


3-Ethylpyrrole was converted to a mixture of Compound A and 2-formyl-4-ethylpyrrole as described in the preparation of Compound A of Example 5. $[M+H]^+ = 124.1$

B. Preparation of 5-ethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine

The mixture obtained above was converted to the title compound in a manner similar to the preparation of Example 5 from compound B of Example 5. Thus, after purification by preparative HPLC, 22 mg (26 %) of 5-ethyl-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine was obtained as white solid. MS: $[M+H]^+ = 269.2$, $[M-H]^- = 267.1$; $^1\text{H NMR}$ (CDCl_3): δ 8.80 (br s, 1H), 7.73 (s, 1H), 7.43 (s, 1H), 6.99-6.89 (m, 2H), 6.60 (s, 1H), 6.55 (d, $J=2.7$ Hz, 1H), 2.45 (s, 1H), 2.87 (q, $J=7.5$ Hz, 2H), 2.00 (s, 3H), 1.31 (t, $J=7.5$ Hz, 3H)

25


Example 17



5-[[2-(Dimethylamino)pyrrolo[2,1-f][1,2,4]triazin-4-yl]amino]-2-methylphenol

To a solution of 4-chloro-2-dimethylaminopyrrolo[2,1-f][1,2,4]triazin-2-amine (50 mg, 0.26 mmol) (*Tetrahedron*, 3037, 52, 1996, José Ma. Quintela, Maria J. Moreira and Carlos Peinador) in ethanol (2.5 mL) under argon was added 5-amino-o-cresol (35 mg, 0.28 mmol). The reaction mixture was stirred overnight at 75°C. Upon cooling, a solid precipitated which was recrystallized from warm ethanol to provide 52 mg (72 %) of the title compound as white solid. MS: $[M+H]^+ = 284.2$, ESI $[M-H]^- = 282.1$; 1H NMR ($CDCl_3$): δ 7.52 (s, 1H), 7.13-6.98 (m, 4H), 6.58 (s, 1H), 3.15 (s, 6H), 2.19 (s, 3H).

Example 18



4-[(3-Hydroxy-4-methylphenyl)amino]-5-phenylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

A. 3-carboethoxy-4-phenylpyrrole

To a 1.0 M solution of lithium hexamethyldisilazide in THF (41 mL, 41 mmol) at -78°C was added dropwise over 45 min. a solution of tosylmethyl isocyanide (8.1 g, 41 mmol) in THF. After the reaction was stirred for an additional 45 min. a solution of trans-ethyl cinnamate (7.3 g, 41 mmol) in THF (40 mL) was added over 40 min. The reaction was warmed to 25°C and stirred for 5 h. The reaction was diluted with ethyl acetate and washed with sat. aqueous $NaHCO_3$. The aqueous layer was extracted three times with ethyl acetate, dried (Na_2SO_4), concentrated and purified by chromatography on silica gel eluting with a gradient of 20-30% ethyl acetate in hexanes to provide 3.4 g (41%) of 3-carboethoxy-4-phenylpyrrole as an off-white solid. 1H NMR ($CDCl_3$): δ 8.62 (br s, 1H), 7.57-7.12 (m, 6H), 6.73 (s, 1H), 4.19 (q, $J=7.1$ Hz, 2H), 1.23 (t, $J=7.1$ Hz, 3H)

B. 3-carboethoxy-4-phenyl-5-formylpyrrole

Compound A was converted to the title compound (74% yield) in a manner similar to the preparation of Compound A of Example 5 from 3-methylpyrrole. $[M-H]^+ = 242$

C. 3-carboethoxy-4-phenyl-5-cyanopyrrole

- 5 To a solution of Compound B (0.55 g, 2.4 mmol) in pyridine (10 mL) under argon at rt was added hydroxylamine hydrochloride (0.18 g, 2.6 mmol). The reaction mixture was stirred for 2 hrs. Acetic anhydride (0.25 mL, 2.6 mmol) was added and the mixture was heated at 95°C for 6 h. Water and ethyl acetate were added to quench the reaction and the mixture was extracted (3 X) with ethyl acetate. The combined
10 extracts were washed with brine, dried (Na_2SO_4) and purified by chromatography on silica gel eluting with a gradient of 25-50% ethyl acetate in hexanes to provide 0.1 g (17%) of Compound C as a solid. ^1H NMR (CDCl_3): δ 9.92 (br s, 1H), 7.52-7.30 (m, 6H), 4.24 (q, $J=7.1$ Hz, 2H), 1.23 (t, $J=7.1$ Hz, 3H)

D. 1-amino-3-carboethoxy-4-phenyl-5-cyanopyrrole

- 15 To a suspension of NaH (60% in oil, 33 mg, 0.83 mmol) in DMF (5 mL) at 0°C was added Compound C (0.1 g, 0.42 mmol) in DMF (3 mL). After 10 min. at 0°C, diphenyl phosphoryl hydroxylamine (0.19 g, 0.83 mmol) was added neat followed by DMF (3 mL). The reaction mixture was stirred for 2 hrs at 25°C and then quenched with pH 7 phosphate buffer (15 mL). The mixture was extracted with ethyl acetate (4
20 x 20 mL). The combined extracts were dried (Na_2SO_4) and purified by chromatography on silica gel eluting with 25-50% ethyl acetate in hexanes to provide 94 mg (85%) of Compound D. ^1H NMR (CDCl_3): δ 10.2 (br s, 1H), 7.55 (s, 1H), 7.47-7.35 (m, 5H), 5.12 (s, 2H), 4.27 (q, $J=7.1$ Hz, 2H), 1.22 (t, $J=7.1$ Hz, 3H)


E. 5-Phenyl-6-carboethoxy-pyrrolo[2,1-f][1,2,4]triazin-4(3H)-one

- 25 To a solution of Compound D (94 mg, 0.4 mmol) in methanol (2 mL) and water (2 mL) was added sodium perborate tetrahydrate (0.28 g, 2 mmol). The reaction mixture was heated to 50°C for 15 hrs and then quenched with water (5 mL). The mixture was extracted with ethyl acetate (3 x 10 mL). The combined organic extracts were washed with brine (10 mL), dried (Na_2SO_4), and concentrated *in vacuo*. The residue was
30 heated with formic acid at 60° C for 5 hrs. The formic acid was removed and the crude material was purified by chromatography silica gel eluting with a gradient of 2-5% methanol in chloroform to provide 26 mg (26%) of Compound E. $[M-H]^+ = 282$

F. 5-phenyl-6-carboethoxy-4-(3-hydroxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine

Phosphorus oxybromide (130 mg, 5 eq) was combined with Compound E (26 mg, 0.092 mmol) and heated to 60°C for 45 min. The melt was poured into ice water and
 5 extracted with ethyl acetate (4 x 5 mL). The extracts were washed with sat. NaHCO₃, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was dissolved in CH₃CN (1.0 mL) and DMF (0.2 mL) and 5-amino-o-cresol (16 mg, 0.13 mmol) was added. The reaction mixture was stirred overnight under argon at 25°C. Solvent was removed *in vacuo*, and the crude material was purified first by radial chromatography
 10 on a 1 mm silica gel plate eluting with a gradient of 2-5% methanol in chloroform, and then by preparative TLC (silica gel, 2% methanol in chloroform) to provide 6 mg (17 %) of the title compound as yellowish solid. [M+H]⁺ = 389; ¹H NMR (CDCl₃): δ 8.05 (s, 1H), 7.98 (s, 1H), 7.47-7.44 (m, 5H), 7.14 (s, 1H), 6.88 (d, *J*=8.2 Hz, 1H), 6.87 (s, 1H), 6.28 (dd, *J*=8.2, 1.6 Hz, 1H), 5.47 (br s, 1H), 4.11 (q, *J*=7.1 Hz, 2H),
 15 2.08 (s, 3H), 1.10 (t, *J*=7.1 Hz, 3H).

Example 19



4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

20 A. 3-methylpyrrole-2,4-dicarboxylic acid dimethyl ester

To a suspension of aluminum chloride (106.4 g, 798 mmol) in dichloroethane (700 mL) at -40°C under nitrogen was added dropwise trichloroacetyl chloride (89 mL, 798 mmol). A solution of 4-methylpyrrole-3-carboxylic acid methyl ester (37 g, 266 mmol, prepared by the procedure analogous to Compound A of Example 18 using
 25 methyl crotonate) in dichloroethane (200 mL) was added and the reaction mixture was gradually warmed to rt. and was stirred over the weekend (65 hr). A cold and pre-prepared aluminum chloride (53.2 g) and trichloroacetyl chloride (44.6 g) in dichloroethane (450 mL) was added to the reaction mixture. After an additional 24 hr, the mixture was carefully poured into an ice-water bath (2 L) and the pH of the

solution was adjusted to 2.0. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The combined organic extract was washed with 3 N HCl, brine, then dried (Na_2SO_4), and concentrated in vacuo to give dark oil. This oil was dissolved in methanol (400 mL), and the resulting solution was cooled 0°C under nitrogen. To this solution was added sodium methoxide (25 % in methanol) until the pH of the solution was 10. After 1 hr, the mixture was concentrated and then diluted with ice water (1 L) and the pH of the mixture was adjusted to 6. The mixture was extracted with dichloromethane (3 x 1 L). The combined extracts were washed with NaHCO_3 , brine, dried (Na_2SO_4) and concentrated in vacuo. The brown solid obtained was purified by chromatography on silica gel eluting with ethyl acetate in hexanes to provide 44.3 g (84%) of Compound A. MS: $[\text{M}+\text{H}]^+ = 196$

B. 1-Amino-3-methylpyrrole-2,4-dicarboxylic acid dimethyl ester

Compound B was prepared from compound A (46 g, 213 mmol) in a manner similar to Compound D of Example 18 except most of the solvent was removed prior to the addition of water. Thus, after purification by chromatography on silica gel eluting with 25-30% ethyl acetate in hexanes, 38 g (84%) of Compound B was obtained as white solid. ESI $[\text{M}+\text{H}]^+ = 213.1$


C. 5-Methylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one-6-carboxylic acid methyl ester

Compound B (38 g, 179 mmol) was combined with formamide (400 mL) and heated to 165°C for 6 hr. The reaction was diluted with water (5 mL), extracted with ethyl acetate (3 x 10 mL), dried (Na_2SO_4) and concentrated. The crude material was purified by washing with ether/hexanes (7/3) to provide 33.4 g (90%) of Compound C as a white solid. ESI MS: $[\text{M}-\text{H}]^- = 206.0$

D. 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared in a manner similar to the preparation of compound F of example 18. Thus, after purification by preparative HPLC, 13.5 mg (42 %) of the title compound was obtained as yellowish solid. ESI $[\text{M}+\text{H}]^+ = 313.2$; ^1H NMR (CDCl_3): δ 7.97 (s, 1H), 7.92 (s, 1H), 7.36 (s, 1H), 7.12 (d, $J=8.1$ Hz, 1H), 6.88 (d, $J=8.1$ Hz, 1H), 3.86 (s, 2H), 2.90 (s, 3H), 2.21 (s, 3H).


Example 20



4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid

A 1.0 M aqueous LiOH solution (0.16 mmol) was added to Example 19 (26 mg, 0.083 mmol) in a mixture of THF (0.6 mL), methanol (0.2 mL), and water (0.2 mL) at 0°C. The reaction mixture was warmed to 25°C and stirred for 2 hrs. An additional solution of LiOH (0.5 mL) was added and the reaction was warmed to 50°C for 1.25 hr. The reaction was brought to pH 7 with 5% aqueous HCl solution and extracted with ethyl acetate (4 x 7 mL). The organic extracts were dried (MgSO₄) and concentrated *in vacuo* to provide 16 mg (67%) of the title compound as off-white solid. MS: [M+H]⁺ = 299.2; ¹H NMR (CD₃OD): δ 7.87 (s, 1H), 7.68 (s, 1H), 7.27 (s, 1H), 6.99 (d, *J*=8.2 Hz, 1H), 6.81 (s, 1H), 2.76 (s, 3H), 2.09 (s, 3H)

Example 21




1-[[4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbonyl]-4-methylpiperazine

To a solution of Example 20 (7 mg, 0.023 mmol) in DMF (0.4 mL) at 25°C was added N-methyl piperazine (3.5 μL, 0.03 mmol), 1-hydroxybenzotriazole (3 mg, 0.023 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (7 mg, 0.05 mmol). After 24 hrs., the mixture was concentrated, diluted with water (2 mL), and extracted with ethyl acetate (5 x 2 mL). The combined organic extracts were washed with saturated aqueous NaHCO₃ (2 x 5 mL). The crude material was purified by chromatography on silica gel eluting with a gradient of 5-10% methanol in chloroform to provide 6.23 mg (70%) of the title compound as white solid. MS: [M+H]⁺=381.3; ¹H NMR (CDCl₃): δ 7.76 (s, 1H), 7.48 (s, 1H), 7.08 (s, 1H), 7.02 (d,

$J=8.2$ Hz, 1H), 6.69 (d, $J=8.2$ Hz, 1H), 3.82-3.44 (m, 2H), 2.54 (s, 3H), 2.53-2.28 (m, 6H), 2.27 (s, 3H), 2.10 (s, 3H)

Example 22




5 **4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide**

Example 20 (7 mg, 0.023 mmol) was converted to 6 mg (65%) of the title compound as white solid in a manner similar to the preparation of Example 21 from Example 20.

MS: $[M+H]^+ = 395.3$; ^1H NMR (DMF- d_7): δ 9.56 (s, 1H), 8.40 (s, 1H), 7.42 (br s,

10 1H), 7.10-7.05 (m, 2H), 3.49-3.47 (m, 12H), 2.49 (s, 3H), 2.14 (s, 3H)

Example 23




4-[(3-Methoxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

15 To a solution of Example 19 (10.8 mg, 0.04 mmol) in CH_3CN (0.5 mL) and methanol (0.1 mL) was added trimethylsilyl diazomethane (21 μL of 2.0 M solution in THF). After 24 hrs. TLC indicated no reaction had occurred. DMF (0.2 mL) and additional trimethylsilyl diazomethane (0.1 mL) were added and the reaction mixture was stirred for additional 24 hrs. After the addition of additional 1.1 eq. of trimethylsilyl


20 diazomethane, LC/MS indicated the disappearance of starting alcohol. Acetic acid (2 drops) was added and the reaction mixture concentrated *in vacuo*. The crude material was purified first by chromatography on silica gel eluting with a gradient of 2-5% methanol in chloroform and then by preparative HPLC to provide 2.8 mg (25%) of

25 Example 23 as white solid. $[M+H]^+ = 327.2$; ^1H NMR (CDCl_3): δ 7.92 (s, 1H), 7.87 (s, 1H), 7.26 (s, 1H), 7.07 (d, $J=8.1$ Hz, 1H), 6.91 (d, $J=8.1$ Hz, 1H), 3.81 (s, 3H), 3.80 (s, 3H), 2.87 (s, 3H), 2.15 (s, 3H).

Example 24

4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-methanol

- 5 To a suspension of Example 19 (23 mg, 0.08 mmol) in CH_2Cl_2 (1 mL) at -78°C under argon was added diisobutyl aluminum hydride (1.0 M in toluene, 0.15 mL). The reaction mixture was gradually warmed to 0°C over 45 min. TLC indicated that starting material remained and an additional 1 equivalent of diisobutyl aluminum hydride was added at 0°C . After 20 min. the reaction mixture was poured into
- 10 aqueous potassium sodium tartrate and stirred for 30 min. The mixture was extracted with CH_2Cl_2 (4 x 5 mL), dried (Na_2SO_4), and purified by chromatography on silica gel eluting with a gradient of 5-10% methanol in chloroform to provide 8 mg (38%) of Example 24 as a solid. MS: $[\text{M}+\text{H}]^+=285.2$; ^1H NMR (CD_3OD): δ 7.70 (s, 1H), 7.56 (s, 1H), 7.19 (s, 1H), 7.08 (d, $J=8.0$ Hz, 1H), 6.89 (d, $J=8.0$ Hz, 1H), 4.68 (s,
- 15 2H), 2.62 (s, 3H), 2.20 (s, 3H)

Example 25

4-[(3-Hydroxy-4-methylphenyl)methylamino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

- 20 A. **5-methyl-6-carbomethoxy-4-(3-tert-butyldimethylsilyloxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine**

To a solution of Example 19 (31 mg, 0.1 mmol) in DMF (1 mL) at 25°C under argon was added t-butyldimethylsilyl chloride (19 mg, 0.13 mmol) and imidazole (11 mg, 0.15 mmol). The reaction mixture was stirred 5 hrs at ambient temperature and then


25 stored at -40°C overnight. Water was added and the mixture was extracted with ethyl

acetate (3 x 5 mL). The organic extracts were washed with brine (10 mL), dried (Na_2SO_4), and purified by chromatography on silica gel eluting with 25% ethyl acetate in hexanes to provide 42 mg (100%) of Compound A. MS: $(\text{M}+\text{H})^+ = 427.4$

B. 5-Methyl-6-carbomethoxy-4-(3-methoxy-4-methylphenyl-N-methylamino)-pyrrolo[2,1-f][1,2,4]triazine

To a solution of Compound A (29 mg, 68 μmol) in THF (0.7 mL) at 0°C was added NaH (5.5 mg, 0.14 mmol). After stirring for 10 min. methyl iodide (17 μL , 0.27 mmol) was added followed by DMF (80 μL) and the reaction mixture was allowed to warm to 25°C over 1 hr. After cooling to 0°C the reaction was quenched with pH 7 phosphate buffer (1 mL) and extracted with ethyl acetate (3 x 2 mL). The combined organic extracts were dried (Na_2SO_4) and purified by chromatography on silica gel eluting with a gradient of 25-50% ethyl acetate in hexanes. The residue was taken up in THF (1 mL) and cooled to 0°C . Tetrabutylammonium fluoride (0.1 mL) was added and the reaction was stirred under argon at 0°C for 45 min. The mixture was quenched with pH 7 phosphate buffer and extracted with ethyl acetate (3 x 2 mL). The organic extracts were dried (Na_2SO_4) and purified by rotary chromatography on a 1 mm silica gel plate eluting with 2% methanol in chloroform to provide 10 mg (50%) of the title compound as a white solid. MS: $[\text{M}+\text{H}]^+ = 327.3$; ^1H NMR (CD_3OD): δ 7.58 (s, 1H), 7.51 (s, 1H), 6.87 (d, $J=7.7$ Hz, 1H), 6.24 (s, 1H), 6.22 (d, $J=7.7$ Hz, 1H), 3.69 (s, 3H), 3.26 (s, 3H), 2.09 (s, 3H), 1.79 (s, 3H)

Example 26



2-Methyl-5-[5-methyl-6-(1H-1,2,4-triazol-1-ylmethyl)pyrrolo[2,1-f][1,2,4]triazin-4-yl]amino]phenol


A. 5-methyl-6-hydroxymethyl-4-(3-tert-butyl dimethylsilyloxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine

To a solution of Compound A of Example 24 (42 mg, 0.1 mmol) in CH_2Cl_2 (1 mL) at -78°C under argon was added diisobutyl aluminum hydride (1.0 M in toluene, 0.30 mL). After 45 min. the reaction mixture was poured into aqueous potassium sodium tartrate and stirred for 40 min at 25°C . The mixture was extracted with ethyl acetate (3 x 5 mL). The combined organic extracts were washed with brine (10 mL), dried (Na_2SO_4) and concentrated to provide compound A which was used directly for the next step without further purification.

B. 5-(1,2,4-pyrazole)methyl-6-hydroxy-4-(3-methoxy-4-methylphenylamino)-pyrrolo[2,1-f][1,2,4]triazine

To a solution of Compound A (20 mg, 50 μmol) in CH_2Cl_2 (1 mL) at 0°C was added triethylamine (75 μmol) and methane sulfonyl chloride (55 μmol). The reaction mixture was warmed to 25°C and stirred for 1 hr. The mixture was concentrated *in vacuo*. In a separate vial 1,2,4-triazole (10 mg, 0.15 mmol) was added to NaH (6 mg, 0.15 mmol) in DMF (1 mL) at 0°C under argon. This mixture was warmed to 25°C and stirred for 15 min. and then cooled back to 0°C . The mesylate was dissolved in 0.5 mL of DMF and added to the second vial. The reaction was warmed to 25°C and stirred for 3 hrs. Water (2 mL) was added and the mixture was extracted with ethyl acetate (3 x 5 mL). The combined organic extracts were dried (Na_2SO_4), and purified by preparative TLC on a 1 mm silica gel plate eluting with 5% methanol in chloroform. The material obtained was dissolved in THF and cooled to 0°C . Tetrabutyl ammonium fluoride (2.0 eq) was added and the mixture was stirred under argon at 0°C for 30 min. The material was concentrated and purified directly by preparative TLC on a 1 mm silica gel plate eluting twice with 5% methanol in chloroform to provide 2.1 mg (10%) of the title compound. MS: $[\text{M}+\text{H}]^+ = 336.2$; ^1H NMR (CD_3OD): δ 8.36 (s, 1H), 7.89 (s, 1H), 7.62 (s, 1H), 7.55 (s, 1H), 7.07 (s, 1H), 6.96 (d, $J=7.9$ Hz, 1H), 6.78 (d, $J=7.9$ Hz, 1H), 5.39 (s, 2H), 2.48 (s, 3H), 2.08 (s, 3H)


Example 27



4-chloro-5-methyl-6-carbomethoxypyrrolo[2,1-f][1,2,4]triazine

Phosphorous oxychloride (2.5 mL) was combined with Compound C of Example 19 (100 mg, 0.483 mmol) and heated at 100°C overnight. The melt was allowed to cool to rt and dissolved in ethyl acetate. The mixture was neutralized with aqueous NaHCO₃ and extracted twice with ethyl acetate. The combined organic washes were dried (Na₂SO₄) and concentrated to provide 101 mg (93%) of Example 27. MS: (M+H)⁺ = 226.6


Example 28



10 4-(2,3-Dihydro-1H-indol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

A mixture of Example 27 (20 mg, 0.09 mmol) and indoline (21 mg, 0.177 mmol) in CH₃CN (1 mL) was shaken for 4 hrs. DMF (0.2 mL) was added, and the crude mixture was purified by preparative HPLC to provide 12.2 mg (45 %) of Example 28 as a white solid. [M+H]⁺=309.2; ¹H NMR (CDCl₃): δ 8.06 (s, 1H), 7.91 (s, 1H), 7.20 (m, 1H), 7.01 (m, 1H), 6.93-6.91 (m, 2H), 4.15 (t, J=7.8 Hz, 2H), 3.81 (s, 3H), 3.09 (t, J=7.8 Hz, 2H), 2.35 (s, 3H).

Example 29




20 4-[(3-Carboxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

A mixture of Example 27 (20 mg, 0.09 mmol), triethylamine (0.1 mL), and 3-amino benzoic acid (24 mg, 0.177 mmol) in CH₃CN (1 mL) was shaken for 4 hrs. The mixture was filtered, washing with CH₃CN, and the crude material was purified by preparative HPLC to provide Example 29 as a solid. [M+H]⁺=327.1; ¹H NMR

(CDCl₃): δ 9.24 (br s, 1H), 8.11 (s, 1H), 7.91 (m, 1H), 7.73-7.58 (m, 4H), 3.80 (s, 3H), 2.83 (s, 3H).

Example 30




5 4-(1H-Indazol-6-ylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

A mixture of Example 27 (20 mg, 0.09 mmol) and 6-aminoindazole (18 mg, 0.13 mmol) in CH₃CN (1 mL) and DMSO (0.5 mL) was shaken for 4 hrs. The mixture was filtered, washed with CH₃CN, and the crude material was purified by preparative

10 HPLC to provide Example 30 as a white solid (13 mg, 45 %). [M+H]⁺=323.1; ¹H NMR (CDCl₃): δ 8.37 (s, 1H), 7.99 (s, 1H), 7.94 (d, *J*=7.4 Hz, 1H), 7.68 (d, *J*=4.2 Hz, 1H), 7.00 (dd, *J*=7.4, 4.2 Hz, 1H), 3.83 (s, 3H), 2.91 (s, 3H).

Example 31




15 4-[[3-Hydroxy-4-[(phenylsulfonyl)amino]phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

A mixture of Example 27 (20 mg, 0.09 mmol) and 4'-amino-2'-hydroxybenzenesulfonamide (18 mg, 0.13 mmol) in DMF (1 mL) was shaken for 4 hrs. The mixture was purified by preparative HPLC to provide Example 31 as a solid

20 (7 mg, 18 %). [M+H]⁺= 454.2; ¹H NMR (CDCl₃): δ 7.92 (s, 1H), 7.76 (s, 1H), 7.69 (d, *J*=7.8 Hz, 1H), 7.51-7.40 (m, 6H), 6.85 (m, 2H), 6.37 (s, 1H), 3.84 (s, 3H), 2.82 (s, 3H).


Example 32



4-(2-Benzothiazolylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

To a 0°C mixture of NaH (60%, 5 mg, 0.106 mmol) in DMF (0.5 mL) was added 2-aminobenzothiazole (16 mg, 0.106 mmol). The reaction mixture was stirred for 10 min. at 0°C. Example 27 (20 mg, 0.09 mmol) was added in DMF (1 mL). The reaction mixture was stirred for 45 min. at 25°C and then quenched with pH 7 phosphate buffer. The mixture was extracted with ethyl acetate. The combined extracts were dried (Na₂SO₄) and purified by preparative HPLC to provide Example 32 as a solid (12 mg, 40%). [M+H]⁺=340.2; ¹H NMR (CDCl₃): δ 8.41 (s, 1H), 8.28 (s, 1H), 7.28 (d, J=7.2 Hz, 1H), 7.08 (m, 2H), 6.57 (d, J=7.2 Hz, 1H), 3.82 (s, 3H), 2.38 (s, 3H).

Example 33




5-Methyl-4-[[3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Phosphorus oxychloride was combined with Example 19 (30 mg, 0.14 mmol) and heated to 100°C for 15 hrs. The melt was cooled and concentrated to remove excess reagent, then extracted with ethyl acetate (20 mL). The extract was washed with saturated aqueous NaHCO₃ (10 mL) dried over MgSO₄ and concentrated *in vacuo* to provide chlorinated compound Example 27. The residue was dissolved in DMF (2 mL) and 3-(methylsulfonylamino)aniline (54 mg, 0.3 mmol) was added. The reaction mixture was stirred for 4 hrs under argon at 25°C. The crude reaction mixture was purified by

preparative HPLC. The material obtained appeared to be a salt of the desired compound. The material was dissolved in ethyl acetate and washed with saturated aqueous NaHCO₃. Evaporation of the solvent gave 24 mg (40 %) of Example 33. MS: [M+H]⁺ = 376.2; ¹H NMR (*d*-DMSO): δ 8.89 (s, 1H), 8.14 (s, 1H), 7.95 (s, 1H), 7.55 (s, 1H), 7.38 (s, 1H), 7.34-7.32 (m, 2H), 7.01 (d, *J*=8.0 Hz, 1H), 3.81 (s, 3H), 3.02 (s, 3H), 2.82 (s, 3H).


Example 34



10 4-[[3-(Hydroxysulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 27 (20 mg, 0.09 mmol) was dissolved in DMF (1 mL) and 3-aminobenzenesulfonic acid (23 mg, 0.13 mmol) was added. After 5 hr at rt, the crude reaction mixture was purified by preparative HPLC. The material obtained appeared to be a salt of the desired compound. The material was dissolved in ethyl acetate and washed with saturated aqueous NaHCO₃. Evaporation of the solvent gave 9.7 mg (30 %) of Example 34. MS: [M+H]⁺ = 363.2; ¹H NMR (CDCl₃/CD₃OD): δ 8.04 (s, 1H), 7.82 (d, *J*=7.1 Hz, 1H), 7.65 (s, 1H), 7.52-7.42 (m, 1H), 7.31-7.26 (m, 2H), 3.73 (s, 3H), 2.75 (s, 3H).

Example 35




20 4-[[3-(Hydrazinocarbonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 35 was prepared from 3-aminobenzhydrazide (30 mg, 0.2 mmol) and Example 27 (30 mg, 1 mmol) in a manner similar to the preparation of Example 28 to

afford 6 mg (18 %) of solid. MS: $[M+H]^+ = 341.2$; ^1H NMR (*d*-DMSO): δ 7.79 (s, 1H), 7.58 (s, 1H), 7.11-7.05 (m, 3H), 6.72 (d, $J = 8.1$ Hz, 1H), 5.23 (br s, 2H), 3.76 (s, 3H), 2.67 (s, 3H)


Example 36



4-[[3-(Aminosulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 27 (20 mg, 0.09 mmol) was treated with 3-aminobenzenesulfonamide (23 mg, 0.13 mmol) in a manner similar to that described for Example 31. Evaporation of the extracting solvent gave 18.6 mg (58 %) of solid. MS: $[M+H]^+ = 362$; ^1H NMR (*d*-DMSO): δ 9.08 (s, 1H), 8.18 (s, 2H), 7.98 (s, 1H), 7.91-7.89 (m, 1H), 7.62-7.58 (m, 2H), 7.42 (s, 2H), 3.81 (s, 3H), 2.84 (s, 3H)


Example 37



4-[[3-[(Butylamino)sulfonyl]phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 27 (20 mg, 0.09 mmol) was treated with *N*-butyl-3-amino-benzenesulfonamide (30 mg, 0.13 mmol) in a manner similar to that described for Example 31. Evaporation of the extracting solvent gave 31 mg (89 %) of solid. MS: $[M+H]^+ = 418.2$; ^1H NMR (CDCl_3): δ 8.07 (s, 1H), 7.97 (s, 1H), 7.85 (s, 1H), 7.84-7.82 (m, 1H), 7.58 (d, $J = 8.0$ Hz, 1H), 7.49-7.45 (m, 2H), 3.82 (s, 3H), 2.93-2.89 (m, 2H), 2.81 (s, 3H), 1.44-1.18 (m, 4H), 0.84 (t, $J = 7.4$ Hz, 3H)


Example 38



4-[(6-Methoxy-3-pyridinyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

Example 27 (20 mg, 0.09 mmol) was dissolved in DMF (1 mL) and 5-amino-2-methoxypyridine (16 mg, 0.13 mmol) was added. The reaction mixture was stirred for 2.5 hrs under argon at 25°C. The crude reaction mixture was concentrated and purified directly by chromatography on silica gel eluting with a gradient of 25-50% ethyl acetate in hexanes. The material obtained was dissolved in ethanol (2 mL) and sodium ethoxide (20%, 0.3 g, 10 eq) was added. The mixture was stirred at 75°C for 3 hrs. The crude material was purified directly by preparative HPLC to provide 4.3 mg (15%) of an off-white solid. MS: $[M+H]^+ = 314.2$; 1H NMR ($CDCl_3$): δ 8.25 (d, $J=2.7$ Hz, 1H), 7.93 (s, 1H), 7.85 (d, $J=8.8$ Hz, 1H), 7.82 (s, 1H), 7.09 (br s, 1H), 6.74 (d, $J=8.8$ Hz, 1H), 3.89 (s, 3H), 3.82 (s, 3H), 2.86 (s, 3H)


Example 39



4-[(3-Cyanophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-aminobenzonitrile (21 mg, 0.177 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 308.2$; 1H NMR ($CDCl_3$): δ 8.23 (s, 1H), 7.97 (s, 1H), 7.93 (s, 1H), 7.75 (d, $J=6.3$ Hz, 1H), 7.46-7.36 (m, 3H), 3.82 (s, 3H), 2.89 (s, 3H)


Example 40



4-[3-(Acetylamino)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3'-
 5 aminoacetanilide (27 mg, 0.177 mmol) by the procedure analogous to Example 28.
 MS: $[M+H]^+ = 340.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.81 (s, 1H), 7.75 (s, 1H), 7.51 (s, 1H),
 7.20-7.07 (m, 3H), 3.80 (s, 3H), 2.84 (s, 3H), 1.90 (s, 3H)


Example 41



4-[(3-Fluoro-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-fluoro-
 4-methylaniline (17 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:
 $[M+H]^+ = 315.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.92 (s, 1H), 7.87 (s, 1H), 7.57 (d, $J=2.8$ Hz,
 15 1H), 7.19-7.09 (m, 3H), 3.81 (s, 3H), 2.87 (s, 3H), 2.20 (s, 3H)

Example 42




5-Methyl-4-[(4-methyl-3-nitrophenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-nitro-
 4-methylaniline (20 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:

$[M+H]^+ = 342.2$; ^1H NMR (CDCl_3): δ 8.39 (s, 1H), 7.96 (s, 1H), 7.91 (s, 1H), 7.30 (d, $J=6.5$ Hz, 1H), 7.19 (d, $J=6.5$ Hz, 1H), 3.83 (s, 3H), 2.89 (s, 3H), 2.54 (s, 3H)

Example 43




5

4-[(3-Cyano-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-cyano-4-methylaniline (18 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:

10 $[M+H]^+ = 323.2$; ^1H NMR (CDCl_3): δ 8.08 (s, 1H), 7.96 (s, 1H), 7.90 (s, 1H), 7.61 (d, $J=8.2$ Hz, 1H), 7.28 (d, $J=8.2$ Hz, 1H), 3.82 (s, 3H), 2.87 (s, 3H), 2.48 (s, 3H)

Example 44




15

4-[(1-Acetyl-2,3-dihydro-1H-indol-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 1-acetyl-6-aminoindoline (23 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:

20 $[M+H]^+ = 366.2$; ^1H NMR (CDCl_3): δ 8.14 (d, $J= 8.6$ Hz, 1H), 7.95 (s, 1H), 7.82 (s, 1H), 7.51 (s, 1H), 7.12 (d, $J= 8.6$ Hz, 1H), 4.03 (t, $J= 7.8$ Hz, 2H), 3.81 (s, 3H), 3.18 (t, $J= 7.8$ Hz, 2H), 2.67 (s, 3H), 2.19 (s, 3H)


Example 45



4-[(3-Amino-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-amino-4-methylaniline (16 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 312.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.92 (s, 1H), 7.90 (s, 1H), 7.11 (s, 1H), 6.96 (d, $J=8.2$ Hz, 1H), 6.74 (d, $J=8.2$ Hz, 1H), 3.81 (s, 3H), 2.85 (s, 2H), 2.18 (s, 3H)

Example 46




4-[(4-Bromo-2-fluorophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 2-fluoro-4-bromoaniline (25.1 mg, 0.13 mmol) by the procedure analogous to Example 28.

MS: $[M+H]^+ = 379.1$; $^1\text{H NMR}$ (CDCl_3): δ 8.52 (t, $J=8.8$ Hz, 1H), 7.95 (s, 1H), 7.92 (s, 1H), 7.60 (br s, 1H), 7.30 (m, 2H), 3.82 (s, 3H), 2.86 (s, 3H)


Example 47



4-(5-isoquinolinylamino)-5-Methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 5-amino isoquinoline (19 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:


$[M+H]^+ = 334.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.12 (s, 1H), 8.09 (s, 1H), 7.10 (t, $J=7.6$ Hz, 1H), 6.83 (d, $J=7.6$ Hz, 1H), 6.70 (d, $J=7.6$ Hz, 1H), 6.59 (dd, $J=7.7, 4.2$ Hz, 1H), 6.01 (d, $J=8.2$ Hz, 1H), 3.82 (s, 3H), 2.71 (s, 3H)

Example 48

4-[(3,4-Dimethylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3,4-dimethylaniline (16 mg, 0.13 mmol) by the procedure analogous to Example 28. MS:

$[M+H]^+ = 310.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.90 (s, 1H), 7.84 (s, 1H), 7.31 (s, 1H), 7.20 (d, $J=8.1$ Hz, 1H), 7.08 (d, $J=8.1$ Hz, 1H), 3.80 (s, 3H), 2.84 (s, 3H), 2.22 (s, 3H), 2.18 (s, 3H)


Example 49

4-[[5-(Hydroxymethyl)-2-methylphenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-amino-5-(hydroxymethyl)-2-methylaniline (18 mg, 0.13 mmol) by the procedure analogous to

Example 28. MS: $[M+H]^+ = 327.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.93 (s, 1H), 7.87 (s, 1H), 7.83 (s, 1H), 7.36-7.21 (m, 3H), 4.65 (s, 2H), 3.80 (s, 3H), 2.87 (s, 3H), 2.28 (s, 3H)


Example 50



4-[(4-Bromo-3-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 4-bromo-3-methylaniline (25 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 376.2$; 1H NMR ($CDCl_3$): δ 7.91 (s, 1H), 7.89 (s, 1H), 7.45 (m, 1H), 7.34 (m, 1H), 7.20 (m, 1H), 3.81 (s, 3H), 2.84 (s, 3H), 2.35 (s, 3H)


Example 51



4-[(4-Chloro-3-nitrophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 4-chloro-3-nitroaniline (23 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 362.7$; 1H NMR ($CDCl_3$): δ 8.48 (s, 1H), 8.00 (s, 1H), 7.94 (s, 1H), 7.75 (dd, $J=8.7, 2.5$ Hz, 1H), 7.48 (d, $J=8.6$ Hz, 1H), 7.38 (s, 1H), 3.83 (s, 3H), 2.89 (s, 3H)


Example 52



4-[(2,3-Dihydro-1,4-benzodioxin-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (2,3-dihydro-1,4-benzodioxin-6-yl)amine (20 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 340.3$; 1H NMR ($CDCl_3$): δ 7.90 (s, 1H), 7.84 (s, 1H), 7.24 (s, 1H), 7.14 (s, 1H), 6.91 (d, $J = 8.4$ Hz, 1H), 6.81 (dd, $J = 8.4, 2.4$ Hz, 1H), 4.20 (m, 4H), 3.80 (s, 3H), 2.83 (s, 3H)


Example 53



5-Methyl-4-[[2-methyl-5-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 2-methyl-5-(methylsulfonyl)aniline (27 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 390.2$; 1H NMR ($CDCl_3$): δ 8.04 (br s, 1H), 7.95 (s, 1H), 7.87 (s, 1H), 7.09-6.99 (m, 2H), 3.82 (s, 3H), 3.00 (s, 3H), 2.89 (s, 3H), 2.28 (s, 3H)

Example 54




5-Methyl-4-[[4-methyl-3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

To a solution of Example 53 (16 mg, 51 μ mol) in pyridine (1 mL) at 0°C was added 4-methyl-3-[(methylsulfonyl)aniline] (4.4 μ L, 87 μ mol). The reaction mixture was stirred for 1 hr at 0°C and then warmed to 25°C and stirred for 4 hrs. Water (5 mL) was added and the mixture was extracted with ethyl acetate (3 x 5 mL). The combined organic extracts were washed with water (10 mL) and brine (10 mL) and dried (Na_2SO_4). The crude material was purified by chromatography on silica gel eluting with 2% methanol in chloroform to provide 6.9 mg (30%) of solid. MS: $[M+H]^+ = 390.2$; 1H NMR ($CDCl_3$): δ 7.93 (s, 1H), 7.86 (s, 1H), 7.77 (s, 1H), 7.35 (d, $J = 8.2$ Hz,

1H), 7.27 (s, 1H), 7.18 (d, $J=8.2$ Hz, 1H), 6.25 (s, 1H), 3.82 (s, 3H), 3.03 (s, 3H), 2.86 (s, 3H), 2.24 (s, 3H)


Example 55



5 4-[(3-Hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-hydroxyaniline (14 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 299.1$; ^1H NMR (CDCl_3): δ 7.93 (s, 1H), 7.88 (s, 1H), 7.70 (s, 1H), 7.30 (br s, 1H), 6.97 (d, $J=8.5$ Hz, 1H), 6.60 (dd, $J=8.4, 2.3$ Hz, 1H), 3.82 (s, 3H), 2.68 (s, 3H).


Example 56



15 4-[[2-(Aminosulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and 3-(aminosulfonyl)aniline (23 mg, 0.13 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 362.1$; ^1H NMR (CDCl_3): δ 8.97 (s, 1H), 8.10 (d, $J=8.1$ Hz, 1H), 7.92 (d, $J=8.1$ Hz, 1H), 7.80 (s, 1H), 7.77 (s, 1H), 7.56 (t, $J=8.2$ Hz, 1H), 7.52 (t, $J=8.2$ Hz, 1H), 5.53 (s, 2H), 3.80 (s, 3H), 2.73 (s, 3H)

Example 57




4-[(3-Hydroxy-4-methoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (28 mg, 0.124 mmol) and 5-amino-2-methoxy-phenol (20 mg, 0.148 mmol) by the procedure analogous to

- 5 Example 28. MS: $[M+H]^+ = 329.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.90 (s, 1H), 7.84 (s, 1H), 7.19 (d, $J = 8.1$ Hz, 1H), 7.18 (s, 1H), 7.02 (d, $J = 8.1$ Hz, 1H), 5.73 (s, 1H), 3.83 (s, 3H), 3.78 (s, 3H), 2.84 (s, 3H).


Example 58



10 **4-[(3-Hydroxy-2-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester**


The title compound was prepared from Example 27 (28 mg, 0.124 mmol) and 3-amino-2-methylphenol (18 mg, 0.148 mmol) by the procedure analogous to Example 28. MS: $[M+H]^+ = 313.2$; $^1\text{H NMR}$ (CD_3OD): δ 7.90 (s, 1H), 7.76 (s, 1H), 6.97 (t, $J = 7.6$ Hz, 1H), 6.80 (d, $J = 7.4$ Hz, 1H), 6.68 (d, $J = 7.4$ Hz, 1H), 3.80 (s, 3H), 2.66 (s, 3H), 2.55 (s, 3H).

Example 59



20 **4-[(2,3-Dihydro-1H-indol-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester**


The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (2,3-dihydro-1H-indol-6-yl)amine (27 mg, 0.13 mmol) by the procedure analogous to Example 29. MS: $[M+H]^+ = 324.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.05 (s, 1H), 7.91 (s, 1H), 6.97 (d, $J = 7.8$ Hz, 1H), 6.31 (s, 1H), 6.23 (d, $J = 7.8$ Hz, 1H), 4.22 (t, $J = 7.7$ Hz, 2H), 3.84 (s, 3H), 2.84 (t, $J = 7.7$ Hz, 2H), 2.55 (s, 3H).

Example 60

5-Methyl-4-[(5-methyl-2-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

- 5 The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (5-methyl-2-pyridinyl)amine (14 mg, 0.13 mmol) by the procedure analogous to Example 30. MS: $[M+H]^+ = 298.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.33 (s, 1H), 8.22 (s, 1H), 6.75 (d, $J=8.7$ Hz, 1H), 6.68 (s, 1H), 6.37 (d, $J=8.7$ Hz, 1H), 3.81 (s, 3H), 2.44 (s, 3H), 1.92 (s, 3H)


10

Example 61

4-[(1H-Benzotriazol-5-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

- The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (1H-Benzotriazol-5-yl)amine (14 mg, 0.13 mmol) by the procedure analogous to Example 30. MS: $[M+H]^+ = 324.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.61 (s, 1H), 7.97 (s, 1H), 7.76 (s, 1H), 7.67 (d, $J=8.1$ Hz, 1H), 7.54 (s, 1H), 7.46 (d, $J=8.1$ Hz, 1H), 3.85 (s, 3H), 2.49 (s, 3H).


20

Example 62

4-[(1,2-Dihydro-4-methyl-2-oxo-7-quinolinyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (1,2-dihydro-4-methyl-2-oxo-7-quinolinyl)amine (14 mg, 0.13 mmol) by the procedure analogous to Example 30. MS: $[M+H]^+ = 314.2$; $^1\text{H NMR}$ (d -DMSO): δ 8.09 (br s, 1H), 7.84 (s, 1H), 7.71 (s, 1H), 7.55 (m, 2H), 7.47 (s, 1H), 3.81 (s, 3H), 2.56 (s, 3H), 2.44 (s, 3H).


Example 63



4-[[3-(Aminosulfonyl)-4-methylphenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and [3-(aminosulfonyl)-4-methylphenyl]amine (14 mg, 0.13 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 376.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.05 (s, 1H), 7.91 (s, 1H), 6.97 (d, $J=7.8$ Hz, 1H), 6.31 (s, 1H), 6.23 (d, $J=7.8$ Hz, 1H), 4.22 (t, $J=7.7$ Hz, 2H), 3.84 (s, 3H), 2.84 (t, $J=7.7$ Hz, 2H), 2.55 (s, 3H).


Example 64



4-[(2,3-Dihydro-1,3-dioxo-1H-isoindol-5-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (20 mg, 0.09 mmol) and (2,3-dihydro-1,3-dioxo-1H-isoindol-5-yl)amine (16 mg, 0.1 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 352.1$; $^1\text{H NMR}$ (CDCl_3): δ 8.38 (s, 1H), 8.18-8.04 (m, 3H), 7.94 (d, $J=7.4$ Hz, 1H), 3.88 (s, 3H), 2.84 (s, 3H).


Example 65



4-[[3-(Hydroxymethyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (18 mg, 0.08 mmol) and [3-(hydroxymethyl)phenyl]amine (12 mg, 0.1 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 313.1$; $^1\text{H NMR}$ (CDCl_3): δ 7.91 (s, 1H), 7.86 (s, 1H), 7.61 (s, 1H), 7.53 (d, $J = 7.9$ Hz, 1H), 7.32 (t, $J = 7.9$ Hz, 1H), 7.10 (d, $J = 7.9$ Hz, 1H), 4.67 (s, 2H), 3.81 (s, 3H), 2.88 (s, 3H).


Example 66



4-(1H-Indol-6-ylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (18 mg, 0.08 mmol) and 1H-indol-6-ylamine (13 mg, 0.1 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 322.1$; $^1\text{H NMR}$ (CD_3OD): δ 8.00 (s, 1H), 7.76 (d, $J = 3.0$ Hz, 2H), 7.74 (s, 1H), 7.59 (d, $J = 8.4$ Hz, 2H), 7.29 (d, $J = 3.0$ Hz, 1H), 6.48 (d, $J = 8.4$ Hz, 1H), 3.83 (s, 3H), 2.72 (s, 3H).


Example 67



4-[(3-Carboxy-4-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (22 mg, 0.1 mmol), 5-aminosalicylic acid (23 mg, 0.15 mmol) and 1 drop of triethylamine by the procedure analogous to Example 31. MS: $[M+H]^+ = 343.1$; 1H NMR (d -DMSO): δ 8.80 (s, 1H), 8.10-8.03 (m, 3H), 7.90 (s, 1H), 7.75 (m, 2H), 6.90 (d, $J=8.9$ Hz, 1H), 3.79 (s, 3H), 2.82 (s, 3H).


Example 68



5-Methyl-4-[(4-phenoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (22 mg, 0.1 mmol) and 5-aminosalicylic acid (28 mg, 0.15 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 322.1$; 1H NMR ($CDCl_3$): δ 7.92 (s, 1H), 7.86 (s, 2H), 7.53 (d, $J=8.4$ Hz, 2H), 7.31-7.26 (m, 3H), 7.19-6.95 (m, 5H), 3.82 (s, 3H), 2.85 (s, 3H).


Example 69



5-Methyl-4-[(1,2,3,4-tetrahydro-1,4-dioxo-6-phthalazinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (22 mg, 0.1 mmol) and (1,2,3,4-tetrahydro-1,4-dioxo-6-phthalazinyl)amine (27 mg, 0.15 mmol) by the procedure analogous to Example 31. MS: $(M-H)^- = 365^-$


Example 70



4-[(6-Methoxy-3-pyridinyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (22 mg, 0.1 mmol) and (1,2,3,4-tetrahydro-1,4-dioxo-6-phthalazinyl)amine (19 mg, 0.15 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 342.2$; 1H NMR ($CDCl_3$): δ 8.25 (s, 1H), 7.94 (s, 1H), 7.85 (dd, $J=8.6, 2.6$ Hz, 1H), 7.82 (s, 1H), 6.79 (d, $J=8.9$ Hz, 1H), 3.89 (s, 3H), 3.82 (s, 3H), 2.84 (s, 3H).


Example 71



5-Methyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (10 mg, 0.044 mmol) and [1-(phenylmethyl)-1H-indazol-5-yl]amine (20 mg) by the procedure analogous to Example 31 to provide 9.1mg (50 %) of solid. MS: $[M+H]^+ = 413.2$; 1H NMR ($CDCl_3$): δ 8.00 (m, 3H), 7.92 (s, 1H), 7.84 (s, 1H), 7.35-7.25 (m, 7H), 5.54 (s, 2H), 3.93 (s, 3H), 2.80 (s, 3H).


Example 72



4-[[6-(Acetylamino)-3-pyridinyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (10 mg, 0.044 mmol) and [6-(acetylamino)-3-pyridinyl]amine (14 mg, 0.09 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 341.2$; $^1\text{H NMR}$ (*d*-DMSO): δ 10.51 (s, 1H), 8.87 (s, 1H), 8.53 (s, 1H), 8.14-8.01 (m, 3H), 7.93 (s, 1H), 3.80 (s, 3H), 2.83 (s, 3H), 2.32 (s, 3H).


Example 73



5-Methyl-4-(1H-pyrazol-3-ylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (10 mg, 0.044 mmol) and 1H-pyrazol-3-ylamine (7 mg, 0.09 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 273.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.94 (d, $J = 7.2$ Hz, 2H), 7.77 (s, 1H), 7.69 (s, 1H), 6.45 (br s, 1H), 3.84 (s, 3H), 2.85 (s, 3H).


Example 74



4-[(4-Methoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (10 mg, 0.044 mmol) and 4-methoxyaniline (11 mg, 0.09 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 313.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.91 (s, 1H), 7.83 (s, 1H), 7.44 (d, $J = 6.9$ Hz, 2H), 6.89 (d, $J = 7.6$ Hz, 2H), 3.81 (s, 3H), 3.76 (s, 3H).


Example 75



4-[(3,4-Dimethoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester


The title compound was prepared from Example 27 (10 mg, 0.044 mmol) and 3,4-dimethoxyaniline (14 mg, 0.09 mmol) by the procedure analogous to Example 31.
 5 MS: $[M+H]^+ = 343.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.91 (s, 1H), 7.85 (s, 1H), 7.21 (s, 1H), 7.00 (dd, $J=8.6, 2.8$ Hz, 1H), 6.82 (d, $J=8.6$ Hz, 1H), 3.84-3.82 (3s, 9H), 2.88 (s, 3H)

Example 76



10 The title compound was prepared from Example 27 (23 mg, 0.10 mmol) and 1,3-dihydroxybenzene (17 mg, 0.15 mmol) by the procedure analogous to Example 32.
 MS: $[M+H]^+ = 300$; $^1\text{H NMR}$ (CD_3OD): δ 8.38 (s, 1H), 7.92 (s, 2H), 7.28 (t, $J=8.2$ Hz, 1H), 6.78-6.70 (m, 2H), 3.86 (s, 3H), 2.82 (s, 3H).

Example 77




15

4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The solution of oxindole (5.32 g, 40 mmol) in THF (150 ml) and DMF (35 ml) was deoxygenated by purging with argon. To this mixture in ice bath, sodium hydride
 20 (60% in oil, 1.7 g, 42 mmol) was added. After 30 min, Example 27 (3.38 g, 15 mmol) was added. After 1 hr. at rt, the resulting mixture was neutralized with acetic acid. The solvent was removed in vacuo. The residue was dissolved in dichloromethane, washed with brine, and dried (MgSO_4). The solution was concentrated to a solid

residue which was triturated with dichloromethane and diethyl ether to afford the title compound as an orange solid (3.5 g, 72%). MS: $[M+H]^+ = 323.1$; $^1\text{H NMR}$ (CDCl_3): δ 7.86 (s, 1H), 7.74 (s, 1H), 7.37 (s, 1H), 7.19 (d, $J=7.6$ Hz, 1H), 7.11 (t, $J=7.6$ Hz, 1H), 7.10 (t, $J=7.6$ Hz, 1H), 7.06 (d, $J=7.6$ Hz, 1H), 3.84 (s, 3H), 2.34 (s, 3H)


5

Example 78

4-(2,3-Dihydro-1-methyl-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid from Example 27 (22 mg, 0.10 mmol) and N-methyloxindole (22 mg, 0.15 mmol) by the procedure analogous to Example 32. MS: $[M+H]^+ = 337.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.90 (s, 1H), 7.40 (s, 1H), 7.16 (m, 2H), 7.02 (t, $J=7.6$ Hz, 1H), 6.90 (d, $J=7.6$ Hz, 1H), 5.28 (s, 1H), 3.86 (s, 3H), 3.36 (s, 3H), 2.37 (s, 3H).


15

Example 79

4-(2,3-Dihydro-2-oxo-1H-benzimidazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a pale yellow solid (12 mg, 37%) from Example 27 (23 mg, 0.1 mmol) and 2,3-Dihydro-2-oxo-1H-benzimidazol (40.2 mg, 0.3 mmol) by the procedure analogous to Example 32. $^1\text{H NMR}$ (CDCl_3): δ 8.37 (s, 1H), 8.32 (s, 1H), 7.95 (br, s, 1H), 7.21-7.10 (m, 4H), 3.88 (s, 3H), 2.51 (s, 3H).


Example 80



5-Methyl-4-(1,2,3,4-tetrahydro-3-oxo-1-quinoxaliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 27 (23 mg, 0.1 mmol) was stirred with 1,2,3,4-tetrahydroquinoxalin-2-one (44.4 mg, 0.3 mmol) in DMF (0.5 mL) for 1 hr at 50°C. Water was added and the resulting solid material was collected, washed with water and dried. The material was triturated with methanol, filtered, and dried again to provide 20 mg (59%) of a white solid. ¹H NMR (*d*-DMSO): δ 8.30-8.25 (m, 2H), 7.12-7.03 (m, 2H), 6.83 (br s, 2H), 4.38 (s, 2H), 3.73 (s, 3H), 2.49 (s, 3H), 1.72 (s, 3H).


Example 81



4-[2,3-Dihydro-5-[(methylsulfonyl)amino]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (24 mg, 0.1 mmol) and (methylsulfonyl)amino]-2-oxo-1H-indole (90 mg, 0.4 mmol) by the procedure analogous to Example 32. After purification directly by preparative HPLC, the desired material was collected, concentrated, and neutralized with aqueous NaHCO₃. The solid was collected and dried to provide a yellow solid (6 mg, 14%). ¹H NMR (CD₃OD): δ 7.99 (s, 1H), 7.69 (s, 1H), 7.04-7.02 (m, 2H), 6.92 (d, *J* = 8.8 Hz, 1H), 3.86 (s, 3H), 2.87 (s, 3H), 2.39 (s, 3H).

Example 82



4-Chloro-5-ethyl pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

A. 4-Ethylpyrrole-3-carboxylic acid ethyl ester

Compound A was prepared from ethyl-trans-2-pentenoate by the procedure analogous to the preparation of Compound A of Example 18.


B. 5-Ethylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one-6-carboxylic acid ethyl ester

- 5 Prepared from Compound A above by the procedure analogous to the preparation of Compound C of Example 19 from Compound A of Example 19.

C. 4-Chloro-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

- Prepared from Compound B by the procedure analogous to the preparation of
10 Example 27. MS: $(M+H)^+ = 254.6$

Example 83




5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

- 15 The title compound (369 mg, 88%) was obtained from Example 82 (290 mg, 1.23 mmol) and 5-amino-2-methoxy pyridine (229 mg, 1.85 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 342.2$; 1H NMR ($CDCl_3$): δ 8.26 (s, 1H), 7.97 (s, 1H), 7.87 (d, $J = 8.4$ Hz, 1H), 7.82 (s, 1H), 6.80 (d, $J = 8.9$ Hz, 1H), 4.28 (q, $J = 7.1$ Hz, 2H), 3.84 (s, 3H), 3.25 (q, $J = 7.6$ Hz, 2H), 1.32 (s, 6H)

20

Example 84




5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid

- To a solution of Example 83 (7 mg, 0.02 mmol) in methanol (2 mL) was added 2.0 M
25 aqueous sodium hydroxide (0.2 mL, 0.4 mmol). The reaction mixture was warmed to 70°C and stirred for 3 hrs. The methanol was removed *in vacuo* and the residue was

brought to pH 1 with 1 N aqueous HCl solution. The mixture was extracted with CH_2Cl_2 . The combined organic extracts were dried (Na_2SO_4) and concentrated *in vacuo* to provide 4 mg (65%) of white solid. MS: $[\text{M}+\text{H}]^+ = 314.1$; ^1H NMR (CDCl_3): δ 8.05 (s, 1H), 8.03 (s, 1H), 7.88 (m, 2H), 6.88 (d, $J = 8.7$ Hz, 1H), 3.90 (s, 1H), 3.26 (q, $J = 7.2$ Hz, 2H), 1.28 (t, $J = 7.2$ Hz, 3H).


Example 85



5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

To a solution of Example 84 (3 mg, 0.01 mmol) in methanol (0.3 mL) and toluene (0.7 mL) was added trimethylsilyl diazomethane (100 μL of 2.0 M solution in THF). After 30 min. acetic acid and methanol were added and the reaction mixture was concentrated *in vacuo*. The crude material was purified by chromatography on silica gel eluting with 20% ethyl acetate in hexanes to provide 3 mg of white solid. MS: $[\text{M}+\text{H}]^+ = 328.1$; ^1H NMR (CDCl_3): δ 8.30 (s, 1H), 7.94-7.90 (m, 3H), 7.01 (s, 1H), 6.81 (d, $J = 8.8$ Hz, 1H), 3.96 (s, 3H), 3.88 (s, 3H), 3.32 (q, $J = 7.6$ Hz, 2H), 1.56 (t, $J = 7.6$ Hz, 3H).

Example 86




[4-[(6-Methoxy-3-pyridinyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester

To a suspension of Example 84 (36 mg, 0.12 mmol) in 1,4-dioxane (1.4 mL) under argon with powdered 4 Å molecular sieves was added triethylamine (19 μL , 0.14 mmol, 1.2 eq) and diphenylphosphoryl azide (30 μL , 0.14 mmol). The mixture was heated at 85°C for 1 hr and then benzyl alcohol (24 μL , 0.23 mmol) was added. The reaction mixture was warmed at 85°C for 15 hrs. The mixture was filtered, concentrated *in vacuo*, and purified directly by rotary chromatography directly on a 1

mm silica gel plate eluting with 2% methanol in chloroform to provide 29 mg (60%) of yellow oil. MS: $[M+H]^+ = 419.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.27 (d, $J=2.7$ Hz, 1H), 7.98 (s, 1H), 7.95 (dd, $J=8.8, 2.7$ Hz, 1H), 7.90 (s, 1H), 7.43-7.35 (m, 5H), 6.81 (d, $J=8.8$ Hz, 1H), 6.46 (br s, 1H), 5.24 (s, 2H), 3.96 (s, 3H), 2.82 (q, $J=7.7$ Hz, 2H), 1.34 (t, $J=7.7$ Hz, 3H).


Example 87



5-Ethyl-4-[(3-hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a solid from Example 82 (24 mg, 0.095 mmol) and 3-hydroxy-4-methylaniline (17 mg, 0.14 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 339.1$; $^1\text{H NMR}$ (CDCl_3): δ 8.00 (s, 1H), 7.59 (s, 1H), 7.10 (s, 1H), 6.88 (d, $J=8.4$ Hz, 1H), 6.52 (d, $J=8.4$ Hz, 1H), 4.33 (q, $J=7.6$ Hz, 2H), 3.20 (m, 2H), 2.11 (s, 3H), 1.34 (t, $J=7.6$ Hz, 3H), 1.11 (t, $J=7.4$ Hz, 3H).


Example 88



5-Ethyl-4-[(4-methoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a solid from Example 82 (15 mg, 0.06 mmol) and 4-methoxy aniline (11 mg, 0.09 mmol) by the procedure analogous to example 31. MS: $[M+H]^+ = 341.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.02 (s, 1H), 7.81 (s, 1H), 7.32 (d, $J=8.9$ Hz, 1H), 6.91 (d, $J=8.8$ Hz, 1H), 4.30 (q, $J=7.1$ Hz, 2H), 3.77 (s, 3H), 3.11 (m, 2H), 1.33 (q, $J=7.1$ Hz, 3H), 1.26 (m, 3H).


Example 89



4-[(3-Amino-4-methylphenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a solid from Example 82 (15 mg, 0.06 mmol) and
 5 1,3-diamino-4-methyl benzene (19 mg, 0.09 mmol) by the procedure analogous to
 example 31. MS: $[M+H]^+ = 340.2$; 1H NMR ($CDCl_3$): δ 7.93 (br s, 2H), 7.91 (s, 1H),
 7.13 (s, 1H), 7.00 (d, $J=6.4$ Hz, 1H), 6.76 (d, $J=6.4$ Hz, 1H), 4.30 (q, $J=7.1$ Hz, 2H),
 3.20 (q, $J=7.6$ Hz, 2H), 2.18 (s, 3H), 1.32-1.23 (m, 6H)

Example 90




10

4-[(6-Chloro-3-pyridinyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a solid from Example 82 (10 mg, 0.044 mmol)
 and 5-amino-2-chloro-pyridine (12 mg, 0.088 mmol) by the procedure analogous to
 15 example 31. MS: ESI $[M+H]^+ = 346.3$; 1H NMR ($CDCl_3$): δ 8.47 (s, 1H), 8.25 (d,
 $J=8.2$ Hz, 1H), 7.98 (s, 1H), 7.90 (s, 1H), 7.31 (d, $J=8.2$ Hz, 1H), 4.28 (q, $J=7.2$ Hz,
 2H); 3.28 (q, $J=7.6$ Hz, 2H), 1.31-1.15 (m, 6H).

Example 91




5-Ethyl-4-(1H-indazol-6-ylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

20

The title compound was obtained as a solid from Example 82 (10 mg, 0.044 mmol) and 6-aminoindazole (10 mg, 0.088 mmol) by the procedure analogous to example 31. MS: $[M+H]^+ = 351.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.39 (s, 1H), 7.97 (m, 3H), 7.68 (d, $J=8.6$ Hz, 1H), 7.46 (s, 1H), 6.98 (d, $J=8.6$ Hz, 1H), 4.29 (q, $J=7.1$ Hz, 2H), 3.30 (q, $J=7.6$ Hz, 2H), 1.34 (t, $J=7.1$ Hz, 3H), 1.31 (t, $J=7.6$ Hz, 3H)


Example 92



5-Ethyl-4-[(4-hydroxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a solid from Example 82 (10 mg, 0.04 mmol) and 4-aminophenol (9 mg, 0.08 mmol) by the procedure analogous to example 31. MS: $[M+H]^+ = 327.2$; $^1\text{H NMR}$ (CDCl_3): δ 7.93 (s, 1H), 7.84 (s, 1H), 7.34 (m, 1H), 7.13 (s, 1H), 6.77 (m, 2H), 4.24 (q, $J=7.2$ Hz, 2H), 3.22 (q, $J=7.6$ Hz, 2H), 1.34-1.25 (m, 6H).


Example 93



5-Ethyl-4-[[4-methyl-3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

To a solution of Example 89 (12 mg, 0.035 mmol) in pyridine (0.35 mL) and CH_2Cl_2 (0.7 mL) at 0°C was added methane sulfonyl chloride (4.8 mg, 0.04 mmol). The reaction mixture was warmed to 25°C and stirred for 4 hrs. CH_2Cl_2 was added and the reaction mixture was washed with aqueous NaHCO_3 and dried (Na_2SO_4). The crude material was purified by chromatography on silica gel eluting with a gradient of 30-50% ethyl acetate in hexanes to provide a solid. MS: $[M+H]^+ = 318.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.03 (s, 1H), 7.96 (s, 1H), 7.91 (d, $J=2.8$ Hz, 1H), 7.38 (d, $J=8.1$ Hz, 1H), 7.28 (dd, $J=8.0, 2.8$ Hz, 1H), 6.35 (s, 1H), 4.31 (q, $J=7.4$ Hz, 3H), 3.38 (q, $J=7.6$ Hz, 2H), 3.21 (s, 3H), 2.38 (s, 3H), 1.40 (m, 6H).


Example 94



4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was obtained as a yellow solid from Example 82 (24 mg, 0.1 mmol) was reacted with oxindole (17 mg, 0.14 mmol) by the procedure analogous to the procedure F of Example 77. MS: $[M+H]^+ = 351.2$; 1H NMR ($CDCl_3$): δ 9.05 (s, 1H), 8.21 (s, 1H), 8.07 (s, 1H), 7.53 (d, $J=8.6$ Hz, 1H), 7.28 (m, 2H), 4.39 (q, $J=7.6$ Hz, 2H), 2.81 (m, 2H), 1.45 (t, $J=7.6$ Hz, 3H), 1.00 (t, $J=7.5$ Hz, 3H).


Example 95



4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid

To a solution of Example 77 (3.3 g, 10.2 mmol) in methanol (600 mL) was added potassium hydroxide (1N aqueous solution, 200 mL) and the mixture was deoxygenated by purging with argon. The reaction mixture was heated to 60°C for 20 hrs. The reaction mixture was cooled and concentrated to about 50 mL and the residue was acidified with concentrated HCl to pH 4. The yellow solid was collected, washed with water, and dried *in vacuo* to afford the title compound (2.9 g, 92%). MS: $[M+H]^+ = 307.1$; 1H NMR (CD_3OD): δ 7.94 (s, 1H), 7.71 (s, 1H), 7.18-7.10 (m, 2H), 6.94-6.86 (m, 2H), 2.45 (s, 3H).

Example 96



4-[(3-Hydroxy-4-methylphenyl)amino]-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

A. [(2-Ethoxy-2-

oxoethyl)(phenylmethyl)amino]methylene]propanedioic acid diethyl ester

5 N-benzylglycine ethyl ester (5.79 g, 30 mmol) was combined with diethyl ethoxymethylene malonate (6.48 g, 30 mmol) and stirred at 120°C for 1 hr. The crude material was used directly for the next reaction.

B. 1-Phenylmethyl-3-hydroxypyrrole-2,4-dicarboxylic acid diethyl ester

10 To a suspension of NaH (60% in oil, washed with hexanes, 500 mg, 12.5 mmol) in toluene (10 mL) was added Compound A (3.63 g, 10 mmol) in toluene (30 mL) dropwise at 50°C. After 2 hr. the mixture was poured into ice water, and acidified with 1 N aqueous HCl. The mixture was extracted three times with ethyl acetate and the combined organic extracts were dried (MgSO₄) and concentrated *in vacuo*. The
15 crude material was purified by chromatography on silica gel eluting with 50% ethyl acetate in hexanes to provide 2.70 g (85%) of Compound B as a pink oil.

C. 1-Phenylmethyl-3-methoxypyrrole-2,4-dicarboxylic acid diethyl ester

Compound B (634 mg, 2 mmol) was stirred in acetone for 10 hrs at rt with methyl
20 iodide (300 mg, 2.1 mmol) and potassium carbonate (500 mg). The mixture was filtered, concentrated, and purified by chromatography on silica gel eluting with 33% ethyl acetate in hexanes to provide 470 mg (71%) of Compound C as a gel.

D. 3-methoxypyrrole-2,4-dicarboxylic acid diethyl

Compound C (27 g, 81.5 mmol) in ethanol (1 L) was mixed with palladium on carbon
25 (10%, 4 g) and ammonium formate (28 g) and hydrogenated at 40 psi at 90°C for 18 hrs. The reaction mixture was cooled to rt, filtered, and concentrated. The crude material (brown oil) was purified by chromatography on silica gel eluting with 25% ethyl acetate in hexanes to provide 13 g (66%) of tan solid.

E. 1-Amino-3-methoxypyrrole-2,4-dicarboxylic acid diethyl ester

30 To a stirred suspension of NaH (60% in oil, 1.76 g, 70 mmol) in DMF (350 mL) under nitrogen at 0° C was added dropwise a solution of Compound D (13 g, 54 mmol) in DMF (200 mL). After 30 min. the mixture was diluted with DMF (750 mL)

and diphenyl phosphoryl hydroxylamine (15.7 g, 67.4 mmol) was added one portion and the reaction mixture was allowed to warm to rt. After 6 hrs the mixture was concentrated and the residue was diluted with water (1 L), extracted with ethyl acetate (3x1 L). The combined organic extracts were dried (MgSO_4), concentrated and
5 purified by chromatography on silica gel eluting with 20% ethyl acetate in hexanes to provide 13 g (93%) of solid.

F. 4-Hydroxy-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

Compound E (100 mg, 0.39 mmol) was combined with formamide (1 mL) and heated
10 at 180°C for 6 hrs. The reaction mixture was cooled to rt. And diluted with water (5 mL). The solid which formed was collected, washed with water, and dried to provide 70 mg (76%) of Compound F.


G. 4-Chloro-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

15 Phosphorous oxychloride (1 mL) was combined with Compound F (23.7 mg, 0.1 mmol) and heated at reflux for 2 hrs. The melt was allowed to cool to rt and phosphorous oxychloride was removed on rotary evaporator. The crude material was used directly for coupling reactions.

H. 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

20 To a solution of Compound G (0.1 mmol, crude) in acetonitrile (2 mL) was added 5-amino-o-cresol (24.6 mg, 0.2 mmol) and the mixture was stirred for 1 hr. The resulting thick slurry was dissolved in methanol and purified by preparative HPLC. The desired fractions were collected, concentrated, and neutralized with aqueous
25 NaHCO_3 . The resulting solid material was collected and dried to provide 11.0 mg (32%) of the title compound. MS: $[\text{M}+\text{H}]^+ = 343$; ^1H NMR (CD_3OD): δ 7.84 (s, 1H), 7.81 (s, 1H), 7.33 (s, 1H), 7.07 (d, $J=8.0$ Hz, 1H), 6.95 (d, $J=8.0$ Hz, 1H), 4.36 (q, $J=7.0$ Hz, 2H), 4.13 (s, 3H), 2.18 (s, 3H), 1.38 (t, $J=7.0$ Hz, 3H).


Example 97



4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

The title compound was prepared from Compound G of Example 96 (0.2 mmol) and
 5 4-bromo-2-fluoro-aniline (76 mg, 0.3 mmol) by the procedure analogous with the
 preparation of Compound H of Example 96. Thus, when the reaction was complete it
 was concentrated and washed with 1 N aqueous HCl to provide a solid which was
 triturated with aqueous NaHCO₃ and water and then dried to provide 58 mg (70.9%)
 white solid. MS: [M+H]⁺ = 409, 411 (1:1); ¹H NMR (*d*-DMSO): δ 8.83 (s, 1H), 7.97-
 10 7.93 (m, 2H), 7.90 (s, 1H), 7.66 (d, *J*=8.4 Hz, 1H), 7.43 (d, *J*=8.4 Hz, 1H), 4.24 (q,
J=7.5 Hz, 2H), 3.97 (s, 3H), 1.26 (t, *J*=7.0 Hz, 3H).

Example 98



4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

A. 4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid


Example 97 (40.9 mg, 0.1 mmol) in methanol (2 mL) was stirred 72 hrs with 2 N
 aqueous NaOH (1 mL). The mixture was then heated at reflux for 2 hrs. The mixture
 20 was concentrated, acidified with 1 N aqueous HCl, and the resulting solid was washed
 with water and dried to provide 38 mg (100%) of solid.

B. 4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

To a solution of Compound A (14 mg, 0.037 mmol) in DMF (0.2 mL) at 25°C was
 25 added 1-(2-aminoethyl)-pyrrolidine (10 mg, 0.088 mmol), 1-hydroxybenzotriazole
 (10 mg, 0.063 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
 hydrochloride (20 mg, 0.105 mmol). The reaction mixture was stirred 24 hrs. The

crude material was purified by preparative HPLC, the product was collected and converted to its HCl salt and lyophilized to provide 9.5 mg (50%) of the title compound. MS: $[M+H]^+ = 477, 479$ (1:1); 1H NMR (CD_3OD): δ 8.33 (s, 1H), 7.86 (s, 1H), 7.73 (d, $J = 9.7$ Hz, 1H), 7.65-7.57 (m, 2H), 4.18 (s, 3H), 3.84-3.78 (m, 4H), 3.48-3.45 (m, 2H), 3.19-3.13 (m, 2H), 2.21-2.05 (m, 4H).


Example 99



4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-methyl-N-[2-(1-pyrrolidiny)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

To a solution of Compound A of Example 98 (20 mg, 0.052 mmol) in DMF (1 mL) was added 1-(2-methylaminoethyl)-pyrrolidine (10 mg, 0.078 mmol) and benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (25 mg, 0.056 mmol). After 1 hr at 25°C the mixture was purified directly by preparative HPLC, the product was collected, converted to its HCl salt and lyophilized to provide 22 mg (80%) of the title compound. MS: $[M+H]^+ = 491, 493$ (1:1); 1H NMR (CD_3OD): δ 8.27 (s, 1H), 7.83 (s, 1H), 7.73 (d, $J = 7.9$ Hz, 1H), 7.65-7.55 (m, 2H), 4.09 (s, 3H), 3.99-3.96 (m, 4H), 3.85 (br s, 2H), 3.58-3.51 (m, 2H), 3.29-3.15 (m, 5H), 2.19-2.07 (m, 4H).

Example 100




5-Methoxy-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

Compound G of Example 96 (0.1 mmol) and 5-amino-2-methoxypyridine (62 mg, 0.5 mmol) in CH_3CN (0.5 mL) was stirred for 2 hrs. The crude mixture was purified by preparative HPLC. The desired material was collected, concentrated, and neutralized with aqueous $NaHCO_3$. The solid was collected, washed with water, and dried to provide 12.5 mg (36%) of red solid. MS: $[M+H]^+ = 344$; 1H NMR ($CDCl_3$): δ 8.30 (s,

1H), 7.95 (d, $J=8.8$ Hz, 1H), 7.78 (d, $J=7.9$ Hz, 2H), 6.77 (d, $J=8.8$ Hz, 1H), 4.30 (q, $J=7.0$ Hz, 2H), 4.11 (s, 3H), 3.89 (s, 3H), 1.33 (t, $J=7.0$ Hz, 3H).


Example 101



5 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

To a suspension of NaH (60%, 44 mg, 1.1 mmol) in THF (1 mL) was added oxindole (133 mg, 1 mmol). The reaction mixture was stirred for 20 min. at rt. Compound G of example 96 (0.1 mmol) was added. The reaction was stirred for 2 hrs at 25°C. The crude material was purified by preparative HPLC followed by chromatography on silica gel eluting with ethyl acetate to provide 5.5 mg (16%) of a yellow solid. MS: $[M+H]^+ = 353$; $^1\text{H NMR}$ (CDCl_3): δ 8.42 (s, 0.4H), 8.10 (s, 0.6H), 7.79 (s, 1H), 7.75-6.88 (m, 4H), 4.33 (m, 2H), 3.57 (s, 3H), 1.37 (m, 3H).


Example 102



15 4-[5-(Aminocarbonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared from Example 27 (10 mg, 0.05 mmol) and 5-aminocarbonylindoline (23 mg, 0.14 mmol) by the procedure analogous to Example 73. MS: $[M+H]^+ = 366.2$; $^1\text{H NMR}$ ($\text{CDCl}_3/\text{CD}_3\text{OH}$): δ 7.92 (s, 1H), 7.76 (s, 1H), 7.48 (d, $J=8.0$ Hz, 1H), 7.25 (s, 1H), 7.05 (d, $J=8.0$ Hz, 1H), 3.82 (s, 3H), 2.25 (s, 3H).


Example 103



5-Methyl-4-[(5-methyl-1H-pyrazol-3-yl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a white solid (2.5 mg, 18%) from Example 27 (11 mg, 0.05 mmol) and 3-amino-5-methylpyrazole (7 mg, 0.07 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 287.2$; ^1H NMR (CDCl_3 with 1 drop of CD_3OD): δ 7.92 (s, 1H), 7.89 (s, 1H), 6.59 (s, 1H), 3.81 (s, 3H), 2.84 (s, 3H), 2.13 (s, 3H).


Example 104



4-(2,3-Dihydro-3-oxo-1H-indazol-2-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a white solid (4.6 mg, 26%) from Example 27 (13 mg, 0.06 mmol) and 3-indazolinone (12 mg, 0.09 mmol) by the procedure analogous to Example 31. MS: $[M+H]^+ = 324.2$; ^1H NMR (CDCl_3): δ 9.65 (s, 1H), 8.14 (s, 1H), 7.85 (s, 1H), 7.48 (d, $J=8.2$ Hz, 1H), 7.42-7.37 (m, 2H), 7.14-7.12 (m, 1H), 3.86 (s, 3H), 2.84 (s, 3H).


Example 105



4-(2,3-Dihydro-3-oxo-1H-indazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a white solid (4.1 mg, 20%) from Example 27 (16 mg, 0.07 mmol) and 3-indazolinone (14 mg, 0.11 mmol) by the procedure analogous to Example 77. $[M+H]^+ = 324$; $^1\text{H NMR}$ (CDCl_3): δ 8.16 (s, 1H), 8.02-7.99 (m, 2H), 7.83-7.81 (m, 1H), 7.58-7.54 (m, 1H), 7.46 (s, 1H), 3.87 (s, 3H), 2.66 (s, 3H)

5


Example 106

4-[(3-Hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-5,6-dicarboxylic acid diethyl ester

The title compound was made from pyrrolo-3,4-dicarboxylic acid diethyl ester in an analogous manner to that of the preparation of Example 13 from 2,4-dimethylpyrrole.

10

MS: $[M+H]^+ = 404.2$; $^1\text{H NMR}$ (CDCl_3): δ 8.03 (s, 1H), 7.69 (s, 1H), 7.47 (s, 1H), 7.08 (d, $J=8.1$ Hz, 1H), 7.06 (d, $J=8.1$ Hz, 1H), 4.34-4.28 (m, 4H), 2.17 (s, 3H), 1.32 (t, $J=7.1$ Hz, 6H).

Example 107


15

4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-5,6-dicarboxylic acid diethyl ester

The title compound was made from pyrrolo-3,4-dicarboxylic acid diethyl ester in an analogous manner to that of the preparation of Example 106 except the last step was carried out as described in Example 77 with a mixture of THF and DMF as the solvent. MS: $[M+H]^+ = 395.2$; $^1\text{H NMR}$ (CD_3OD): δ 7.83 (s, 1H), 7.76 (s, 1H), 7.55 (d, $J=8.5$ Hz, 1H), 7.18-7.16 (m, 1H), 7.01-6.76 (m, 2H), 4.24 (q, $J=7.1$ Hz, 2H), 3.50 (q, $J=7.2$ Hz, 2H), 1.30 (t, $J=7.1$ Hz, 3H), 0.86 (t, $J=7.2$ Hz, 3H).

20

Example 108



[4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester

A. 5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid

phenylmethyl ester


To a solution of Example 20 (11.5 mg, 60 μ mol) in 1,4-dioxane (0.6 mL) under argon with powdered 4 Å molecular sieves was added triethylamine (10 μ L, 71 μ mol), diphenylphosphoryl azide (15 μ L, 71 μ mol) and benzyl alcohol (12 μ L, 0.12 mmol). The reaction was warmed at 50°C for 15 hrs. The mixture was concentrated *in vacuo* and chromatographed directly on silica gel eluting with a gradient of 2-5% methanol in chloroform to provide 8 mg (50%) of white solid. $(M+H)^+ = 299.2$

B. [4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester

Phosphorous oxybromide (5 eq.) was combined with Compound A (16 mg, 0.054 mmol) and heated to 60°C for 20 min. The melt was poured into ice water and extracted with ethyl acetate (4 x 5 mL). The extracts were washed with aqueous NaHCO_3 , dried (Na_2SO_4) and concentrated *in vacuo*. The residue was dissolved in a mixture of CH_3CN (0.5 mL) and DMF (0.1 mL) and 5-amino-o-cresol (10 mg, 0.081 mmol) was added. The reaction mixture was stirred overnight under argon at 25°C.

Solvent was removed *in vacuo*, and the crude material was purified by rotary chromatography on a 1 mm silica gel plate eluting with 2% methanol in chloroform to provide 3.8 mg (20%) of white solid. MS: $[M+H]^+ = 404.2$; ^1H NMR (CD_3OD): δ 7.79 (s, 1H), 7.70 (s, 1H), 7.44-7.31 (m, 5H), 7.16 (s, 1H), 7.05 (d, $J=7.7$ Hz, 1H), 6.87 (d, $J=7.7$ Hz, 1H), 5.19 (s, 2H), 2.48 (s, 3H), 2.16 (s, 3H).


Example 109



[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester

Example 95 (29 mg, 0.09 mmol) was converted to the title compound in an analogous manner to the preparation of Example 108 to afford the title compound as yellow oil (5 mg, 13%). MS: $[M+H]^+ = 414$; 1H NMR ($CDCl_3$): δ 7.94 (s, 1H), 7.82 (s, 1H), 7.41-7.34 (m, 5H), 7.17-7.14 (m, 1H), 7.04-7.02 (m, 1H), 6.93-6.90 (m, 2H), 6.44 (s, 1H), 5.23 (s, 2H), 2.12 (s, 3H).


Example 110



4-(5-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (9 mg, 28%) from Example 27 (20 mg, 0.09 mmol) and 5-fluorooxindole (27 mg, 0.18 mmol) by the procedure analogous to that of Example 32. MS: $[M+H]^+ = 341.2$; 1H NMR ($CDCl_3$): δ 7.99 (s, 1H), 7.92 (br s, 1H), 7.48 (s, 1H), 6.87 (m, 3H), 3.91 (s, 3H), 2.44 (s, 3H).


Example 111



4-[5-(Aminosulfonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (10 mg, 28%) from Example 27 (20 mg, 0.09 mmol) and 5-aminosulfonyloxindole (38 mg, 0.18 mmol) by the procedure analogous to Example 32. MS: $[M-H]^- = 400.1$; 1H NMR (CD_3OD): δ 8.05 (s, 1H), 7.83 (s, 1H), 7.65 (d, $J=8.2$ Hz, 1H), 7.07 (d, $J=8.3$ Hz, 2H), 3.84 (s, 3H), 2.28 (s, 3H).


Example 112



4-[2,3-Dihydro-5-[[[2-(4-morpholinyl)ethyl]amino]sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (8 mg, 10%) from Example 27 (20 mg, 0.09 mmol) and 5-(2-morpholinylethylamino)sulfonyloxindole (59 mg, 0.18 mmol) by the procedure analogous to Example 32. MS: $[M+H]^+ = 515.2$; 1H NMR (CD_3OD): δ 8.07 (s, 1H), 7.79 (s, 1H), 7.62 (m, 2H), 7.14 (d, $J=8.2$ Hz, 1H), 4.20 (m, 2H), 3.84 (s, 3H), 3.80 (m, 2H), 3.30-3.12 (m, 8H), 2.30 (s, 3H).

Example 113




10

4-[2,3-Dihydro-2-oxo-5-[[[2-(1-pyrrolidinyl)ethyl]amino]sulfonyl]-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (12 mg, 52%) from Example 27 (10 mg, 0.05 mmol) and 5-(2-pyrrolidinylethylamino)sulfonyloxindole (27 mg, 0.09 mmol) by the procedure analogous to Example 32. MS: $[M+H]^+ = 499.2$; 1H NMR (CD_3OD): δ 8.06 (s, 1H), 7.79 (s, 1H), 7.64 (m, 2H), 7.14 (d, $J=8.1$ Hz, 1H), 3.86 (s, 3H), 3.74 (m, 2H), 3.33 (m, 4H), 3.30 (m, 2H), 2.40 (s, 3H), 2.17-2.24 (m, 4H).

15

Example 114




20

4-[2,3-Dihydro-5-(4-morpholinylsulfonyl)-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (10 mg, 47%) from Example 27 (10 mg, 0.05 mmol) and 5-morpholinylsulfonyloxindole (26 mg, 0.09 mmol) by the procedure analogous to Example 32. MS: $[M+H]^+ = 472.2$; 1H NMR (CD_3OD): δ

8.04 (s, 1H), 7.88 (m, 2H), 7.66 (d, $J=7.8$ Hz, 1H), 7.14 (d, $J=7.9$ Hz, 1H), 3.88 (s, 3H), 3.74-3.70 (m, 4H), 3.36-3.30 (m, 4H), 2.41 (s, 3H).


Example 115



5 4-[2,3-Dihydro-5-[(2-hydroxyethyl)amino]sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (14 mg, 60% overall) from Example 27 (10 mg, 0.05 mmol) and 5-(2-tert-butyldimethylsilyloxy ethyl)sulfonyloxindole (34 mg, 0.1 mmol) by the procedure analogous to Example 32
 10 followed by desilylation by tetrabutylammonium fluoride. MS: $[M+H]^+ = 446.2$; 1H NMR ($CDCl_3/CD_3OH$): δ 7.84 (s, 1H), 7.55 (s, 1H), 7.28 (d, $J=8.6$ Hz, 1H), 7.20 (s, 1H), 6.78 (d, $J=8.6$ Hz, 1H), 3.78 (s, 3H), 3.60 (t, $J=7.4$ Hz, 2H), 2.77 (t, $J=7.4$ Hz, 2H), 2.20 (s, 3H).

Example 116




15

4-[5-[(Dimethylamino)sulfonyl]-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (9 mg, 47%) from Example 27 (10 mg, 0.05 mmol) and 5-dimethylaminosulfonyl oxindole (22 mg, 0.09 mmol) by the
 20 procedure analogous to Example 32. MS: $[M+H]^+ = 430$. 1H NMR ($CDCl_3$): δ 9.56 (s, 1H), 8.05 (s, 1H), 7.65-7.54 (m, 3H), 7.15 (d, $J=8.2$ Hz, 1H), 3.87 (s, 3H), 2.73 (s, 6H), 2.43 (3H, s).


Example 117



4-[2,3-Dihydro-5-[(methylamino)sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (2 mg, 10%) from Example 27 (10 mg, 0.05 mmol) and 5-methylaminosulfonyl oxindole (22 mg, 0.09 mmol) by the procedure analogous to Example 32. MS: $[M-H]^- = 414$. 1H NMR ($CDCl_3/CD_3OH$): δ 7.98 (s, 1H), 7.76 (s, 1H), 7.54 (s, 1H), 7.31 (d, $J=7.6$ Hz, 1H), 6.85 (d, $J=7.6$ Hz, 1H), 3.82 (s, 3H), 2.28 (s, 3H), 2.21 (s, 3H).


Example 118



4-(5-Cyano-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

The title compound was prepared as a yellow solid (10 mg, 65%) from Example 27 (10 mg, 0.05 mmol) and 5-cyanooxindole (16 mg, 0.1 mmol) by the procedure analogous to Example 32. ESI $[M+H]^+ = 348.2$; 1H NMR (d -DMSO): δ 8.20 (s, 1H), 7.91 (s, 1H), 7.48 (d, $J=8.1$ Hz, 1H), 7.24 (s, 1H), 7.02 (d, $J=8.1$ Hz, 1H), 3.81 (s, 3H), 2.24 (s, 3H).

Example 119




4-(2,3-Dihydro-6-methyl-2-oxo-1H-pyrazolo[2,3-d]pyrimidin-3-yl)-5-

6-methyl-5,7-diazaoxindole

To a solution of ethyl (4-amino-2-methylpyrimidin-5-yl) acetate (WO 99/10349, 0.975 g, 5 mmol) in THF (30 ml), was slowly added potassium *t*-butoxide (1 M in THF, 5 mL). After one hour, the mixture was neutralized with acetic acid to pH 5. The volatiles were removed in vacuo and the residue was purified by flash column chromatography (silica gel, 5-8 % MeOH in dichloromethane) to afford a yellow solid (680 mg, 91%).


B) Methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid**methyl ester**

To a solution of 6-methyl-5,7-diazaoxindole (67 mg, 0.45 mmol) in DMF (2 ml) and THF (1 ml) was added sodium hydride (60% in oil, 20 mg, 0.5 mmol). After stirring for 20 min, Example 27 (34 mg, 0.15 mmol) was added and the mixture was stirred at rt overnight. The mixture was neutralized with acetic acid. Dichloromethane (10 ml) was added to the mixture and the resulting precipitate was collected and washed with small amount of dichloromethane, water, and dried in vacuo to give orange solid (32 mg, 63%). MS: (M+H) = 359

Example 120

4-(2,3-Dihydro-2-oxo-1H-pyrazolo[2,3-b]pyridin-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

To a solution of 7-azaoxindole (Tetrahedron.Lett. 1987, 28, 4027) (60 mg, 0.45 mmol) in DMF (2 mL) and THF (1 mL) was added sodium hydride (60% in oil, 20 mg, 0.5 mmol). After stirring for 20 min, Example 27 (34 mg, 0.15 mmol) was added and the mixture was stirred at rt overnight. The solution was neutralized with acetic acid and dichloromethane (10 ml) was added to the mixture. The resulting solid was collected, washed with small amount of dichloromethane and water, and dried in vacuo to give a yellow solid (35 mg, 72%). LC-MS: (M+H)⁺ = 324.

Example 121

4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

A) 4-Chloro-6-carbomethoxypyrrolo[2,1-f][1,2,4]triazine

Prepared according to the procedure described for Example 27 except using 2-methoxycarbonyl pyrrole as the starting pyrrole.


B) 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Compound A was converted to the title compound using a procedure similar to that used for the preparation of Example 120.

5


Examples 122 to 125 were prepared by a procedure analogous to the preparation of Example 120 using appropriate reagents known in the literature (WO 97/42187).

Example 122




10 4-[6-Fluoro-2-hydroxy-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 123




15 4-[6-Bromo-2-hydroxy-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 124




20 4-[2,3-Dihydro-2-oxo-6-(trifluoromethyl)-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 125



4-[2,3-Dihydro-6-(methylsulfonyl)-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Example 126



5 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

To a solution of Example 95 (50 mg, 0.16 mmol) in DMF (1 mL), dichloromethane (0.5 mL) were added PyBrop (113 mg, 0.24 mmol) and diisopropylethyl amine (0.08 mL, 0.5 mL). After 10 min, 1-(3-aminopropyl)pyrrolidine (61 mg, 0.48 mmol) was added. After 15 h, the reaction mixture was purified by preparative RP HPLC. The yellow oil obtained was converted to the HCl salt and lyophilized to afford a red orange solid (25 mg, 34%). MS: (M+H)⁺ = 419.


The compounds named in Example 127 were prepared from Example 95 and appropriate amines by using the procedure described for Example 126.

15 Example 127

- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 20 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[2-(4-morpholinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-methyl-1-piperazinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 25 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1H-1,2,3-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(2H-1,2,3-triazol-2-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;

- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1H-1,2,4-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(2-methyl-1H-imidazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 5 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[4-(4-morpholinyl)butyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 10 carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;

Example 128




- 15 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-N-[3-(1H-1,2,4-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide
 A) 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid

Example 101 was hydrolyzed by treatment with aqueous 1N KOH in methanol at 55
 20 °C for 2 h. The reaction mixture was adjusted to pH = 3 by addition of aqueous HCl. Partial concentration in vacuo precipitated a yellow solid which was filtered and triturated with water followed by ether to afford the acid (80% yield). MS: (M+H)⁺ = 325.

- B) 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-N-[3-(1H-1,2,4-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide
 25 Procedure same as that for Example 126 except N-[3-(1H-1,2,4-triazol-1-yl)propylamine was used. MS: (M+H)⁺ = 433.

Example 129




4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

Compound A of Example 128 was converted to the title compound by the procedure same as that for Example 126 except N-[3-(4-morpholinyl)propylamine was used.

MS: (M+H)⁺ = 451.

Example 130



1,3-Dihydro-3-[5-methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one

A) 4-Phenoxy-5-methyl-6-carbomethoxypyrrolo[2,1-f][1,2,4]triazine

To a solution of phenol (705 mg, 7.5 mmol) in a mixture of THF (10 mL) and DMF (10 mL), was added NaH (60% in oil, 300 mg, 7.5 mmol). After 30 min, Example 27 (675 mg, 3.0 mmol) was added. After 1 h, the solvent was removed and the residue was poured into 5% aqueous K₂CO₃ solution. The precipitate was collected, washed with water, and dried in vacuo to afford Compound A as white solid (800 mg, 94%). MS: (M+H)⁺ = 284

B) 4-Phenoxy-5-methyl-6-hydroxymethylpyrrolo[2,1-f][1,2,4]triazine

To a solution of Compound A (700 mg, 2.47 mmol) in toluene (20 mL) at -60 °C, was added DIBAL (1.5 M in toluene, 6 mmol). After stirring at 0 °C for 1 h, aqueous 1N HCl (30 mL) was added and the mixture was stirred for 30 min. The mixture was then diluted with dichloromethane (DCM). The organic layer was separated, dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography (silica gel, 2 % MeOH in DCM) to afford Compound B as a solid (610 mg, 96%). MS: (M+H)⁺ = 256

C) 4-Phenoxy-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxaldehyde

A mixture of Compound B (500 mg, 1.96 mmol) and MnO₂ (3.0 g) in toluene (30 mL) was heated at 60 °C for 1 h. After cooling to rt, the mixture was filtered through a pad of silica gel and washed with ethyl acetate. After concentration in vacuo, Compound C was obtained as a white solid (420 mg, 85%). MS: (M+H)⁺ = 254

5 **D) 4-Phenoxy-5-methyl-6-hydroxypyrrolo[2,1-f] [1,2,4]triazine**

A mixture of Compound C (708 mg, 2.8 mmol) and m-CPBA (55-85% pure, 800 mg) in dichloroethane (50 mL) was stirred at rt for 15 hrs. Another portion of m-CPBA (250 mg) was added. After 4 h, the mixture was diluted with DCM and washed with aqueous NaHCO₃ solution. The organic layer was concentrated and the residue was
10 diluted with MeOH and stirred with K₂CO₃ (250 mg) for 1 h. The mixture was concentrated and the residue was diluted with DCM, washed with 2 % aqueous citric acid, dried (MgSO₄). The product was purified by flash column chromatography (4% MeOH in DCM) to afford Compound D (245 mg, 36%) as white solid and Compound C (300 mg, 42%) was recovered. MS: (M+H)⁺ = 242

15 **E) 4-Phenoxy-5-methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f] [1,2,4]triazine**

The solution of Compound D (40 mg, 0.16 mmol) in a mixture of DMF (2 mL) and THF (1 mL) was treated with NaH (60 % in oil, 0.18 mmol). After 20 min, 2-(1H-1,2,-triazol-1-yl)ethyl mesylate (80 mg, 0.27 mmol) was added. The mixture was
20 stirred at rt for 1 h, and at 80 °C for 2 hr. The mixture was then cooled, diluted with DCM, washed with aqueous NaH₂PO₄ solution, dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography (silica gel, 100% DCM to 5% MeOH in DCM) to yield Compound E (17 mg, 32%) as white solid. MS: (M+H)⁺ = 337

25 **F) 5-Methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f] [1,2,4]triazin-4(3H)-one**

A mixture of Compound E (15 mg, 0.045 mmol) in ethanol (10 mL) and HCl (1N, 5 mL) was heated at 80 °C in a sealed tube for 4 h. The mixture was cooled and the volatiles were removed in vacuo. The residue was purified by flash column
30 chromatography (5% MeOH in DCM) to give a white solid (10 mg, 86%). MS: (M+H)⁺ = 261.

G) 4-Chloro-5-methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f] [1,2,4]triazine


A mixture of Compound F (10 mg, 0.04 mmol) and POCl₃ and DIPEA (8 mL) was heated at 90 °C in a sealed tube for 1 h. The volatiles were removed in vacuo. The residue was diluted with DCM, washed with ice-cold aqueous NaHCO₃ solution, dried (MgSO₄) and concentrated. After removal of solvent, Compound G was
 5 obtained as a yellow solid 10 (10 mg, 100%). It was used without further purification.

H) 1,3-Dihydro-3-[5-methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one

A solution of oxindole (65 mg, 0.5 mmol) in DMF (2 mL) was purged with argon.
 10 NaH (60% in oil, 20 mg, 0.5 mmol) followed by Compound G (10 mg, 0.04 mmol) were added to the reaction mixture. After 2 h, acetic acid (50 mL) was added to quench the reaction. The volatiles were removed in vacuo and the residue was purified by flash column chromatography (silica gel, 5% MeOH in DCM) to give the title compound as an orange solid (6 mg, 42%). MS: (M+H)⁺ = 376.


15

Example 131



Procedure described for Example 130 was used except in Step E, iodomethane was used and the reaction mixture was stirred at rt. MS: (M+H)⁺ = 295.

Example 132



20

4-(3-Hydroxy-4-methoxyphenyl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

A) 5-Methylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one-6-carboxylic acid

To a solution of the Compound C of Example 19 (1.035 g, 5.00 mmol) in a mixture of
 25 tetrahydrofuran/methanol/water (50 mL, 3:1:1) was added lithium hydroxide (2.062 g, 49.1 mmol). The reaction mixture was stirred at 55 °C for 12 h, then cooled to 0 °C

and neutralized by 3N HCl. The organic solvents were removed and the aqueous solution brought to pH 4 with 1 N HCl. The resulting precipitate was filtered, rinsed with cold water and air dried to afford the Compound A as an off-white solid (0.965 g, 100%).

5 **B) 4-Chloro-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide**

A suspension of Compound A (2.00 g, 10.4 mmol) in phosphorous oxychloride (8 mL) was stirred at 100 °C over 4 h. The solvent was removed *in vacuo* using toluene to assist in the removal. The resulting green solid was suspended in acetonitrile (20 mL) at 0 °C and treated with sufficient triethylamine (5 mL) to bring the solution to pH 10. 4-(3-aminopropyl)morpholine (1.5 mL, 10.3 mmol) was added and the solution was allowed to stir at ambient temperature over 1 h. The reaction mixture was poured into saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was dried (MgSO₄) and the volatiles were removed *in vacuo* to afford Compound B as a yellow solid (1.75 g, 50%). It was used without further purification.


C) 4-(3-Hydroxy-4-methoxyphenyl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

To a solution of Compound B (25.5 mg, 0.075 mmol) in DMF (1 mL) was added 5-amino-2-methoxyphenol (21 mg, 0.15 mmol). After 4 h at rt the volatiles were removed *in vacuo*. Chromatography on silica gel eluting with a gradient of 2 to 10% methanol in dichloromethane yielded a white solid. This material was suspended in acetonitrile (1 mL) and dichloromethane (1 mL) and treated with 1 N hydrogen chloride in diethyl ether to afford the HCl salt of the title compound as a grey solid (21 mg, 58%). MS: (M+H)⁺ = 441.

Example 133

The following two compounds were prepared by treatment of compound B of Example 132 and appropriate amine using the procedure described for the synthesis of Example 132.

30 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
4-[(4-Bromophenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide.

Example 134

5 **4-(5-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide**


To a solution of 5-fluoro oxindole (36 mg, 0.24 mmol) in DMF (1 mL) was added NaH (5.9 mg, 0.23 mmol). After 30 min at rt, a solution of Compound B of Example 132 (24 mg, 0.072 mmol) in DMF (1 mL) was added and the resulting mixture was stirred at ambient temperature over 1 h. The solvent was removed *in vacuo* and the mixture was purified by RP HPLC. The methanol in the desired HPLC fractions was removed *in vacuo* and the resulting aqueous solution neutralized using saturated sodium bicarbonate solution, then extracted with ethyl acetate. The organic layer was dried (MgSO₄) and the volatiles were removed *in vacuo*. The solid obtained was dissolved in acetonitrile/MeOH and treated with 1 N HCl in diethyl ether. The mixture was stirred at ambient temperature over 1 h and the solvents removed *in vacuo*. The HCl salt of the title compound was obtained as an orange solid (18 mg, 51%). MS: (M+H)⁺ = 453.

Example 135

20 The following two compounds were prepared by treatment of compound B of Example 132 with appropriately substituted oxindole using the procedure described for the synthesis of Example 134.

- 4-(6-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 4-[5-(Aminosulfonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide.

Example 136



4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propanoic acid methyl ester

A) 4-Phenoxy-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propenoic acid methyl ester.

DBU (1.42 mL, 9.49 mmol) was added to a solution of the Compound C of Example 130 (600 mg, 2.37 mmol) and methyl diethylphosphonoacetate (1.74 mL, 9.49 mmol) in 1,2-dichloroethane (20 mL). After stirring at rt overnight, the reaction mixture was diluted with dichloromethane and washed with aqueous 2% citric acid, brine, dried (MgSO₄), and concentrated. The organic extract was concentrated and the residue was purified by chromatography on silica gel and elution with 20% ethyl acetate (EtOAc)/DCM to afford a white solid (710 mg, 97%). MS: (M+H)⁺ = 310.

B) 4-Hydroxy-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propanoic acid methyl ester.

Pd/C (10%, 70 mg) was added to a solution of the Compound A (710 mg, 2.30 mmol) in a solvent mixture EtOAc/MeOH/THF/AcOH (100 mL/100 mL/20 mL/2 mL). The suspension was stirred under hydrogen for 2h. The reaction mixture was passed through Celite, the Celite was washed with MeOH and the filtrate was concentrated in vacuo to give crude product. Trituration with hexanes afforded Compound B as a white solid (430 mg, 88%). MS: (M+H)⁺ = 236.

C) 4-Chloro-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propanoic acid methyl ester.


A mixture of diisopropylethylamine (0.24 mL, 1.4 mmol), Compound B (220 mg, 0.94 mmol) and POCl₃ (3 mL) was heated in a sealed bottle at 80 °C. After 2 h, the mixture was cooled down to rt and concentrated in vacuo to give a residue. The residue was partitioned between DCM and aqueous NaHCO₃ solution. The DCM layer was separated, dried (MgSO₄) and concentrated in vacuo to give a dark green solid. Purification by chromatography on silica gel and elution with 20% EtOAc/DCM afforded yellow solid (220 mg, 92%). MS: (M+H)⁺ = 254.

D) 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propanoic acid methyl ester.

NaH (60% in oil, 28 mg, 0.71 mmol) was added to a solution of oxindole (94 mg, 0.71 mmol) in DMF (2 mL) under argon and the mixture was stirred for 10 min.

- 5 Compound C (60 mg, 0.24 mmol) was added to the solution. After 1 h at rt, the reaction was quenched by the addition of acetic acid and diluted with DCM. The organic solution was washed with water, dried (MgSO₄), and concentrated in vacuo to give crude product. Purification by chromatography on silica gel and elution with 20% EtOAc/DCM afforded the title compound as a pure yellow solid (78 mg, 94%).
- 10 MS: (M+H)⁺ = 351.

Example 137



1,3-Dihydro-3-[5-methoxy-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one

- 15 **A) 4-Hydroxy-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-methanol**
Compound F of Example 96 (3.56 g, 15 mmol) was combined with lithium tri-*tert*-butoxyaluminumhydride (1 M solution in THF, 60 mL, 60 mmol) and heated at reflux overnight. The reaction mixture was allowed to cool to rt and quenched with 1 N aqueous HCl. The mixture was concentrated to remove volatiles and the remaining
- 20 material was combined with 100 g of silica gel and applied to a flash silica gel column which was eluted with ethyl acetate to provide 2.65 g (90%) of compound A. MS: [M+H]⁺ = 196.

B) 2,2-Dimethylpropanoic acid [6-(hydroxymethyl)-5-methoxy-4-oxopyrrolo[2,1-f][1,2,4]triazin-3(4H)-yl]methyl ester


- 25 Compound A (195 mg, 1 mmol) was dissolved in 1.5 mL of N,N-dimethylformamide. Sodium hydride (60% in oil, 48 mg, 1.2 mmol) was added and the reaction mixture was stirred at rt for 0.5 hr. Chloromethyl pivalate (181 mg, 1.2 mmol) was added and the mixture was stirred for 1 hr. Water was added and the mixture was extracted with ethyl acetate (3 x 10 mL). The combined extracts were dried (Na₂SO₄), concentrated
- 30 *in vacuo* and purified by flash column chromatography on silica gel eluting with 33%

ethyl acetate in hexanes to provide 260 mg (84%) of compound B as a solid. MS: $[M+H]^+ = 310$.

C) 2,2-Dimethylpropanoic acid [6-formyl-5-methoxy-4-oxopyrrolo[2,1-f][1,2,4]triazin-3(4H)-yl]methyl ester

- 5 Compound B (740 mg, 2.39 mmol) was suspended in toluene (10 mL) with manganese dioxide (835 mg, 9.6 mmol) and heated at 100°C for 3 hr. The reaction mixture was cooled to rt, filtered, and the precipitate was washed with ethyl acetate. The filtrate was concentrated *in vacuo* to provide 660 mg (90%) of compound C as a solid. MS: $[M+H]^+ = 308$.

10 **D) 2,2-Dimethylpropanoic acid [6-formyloxy-5-methoxy-4-oxopyrrolo[2,1-f][1,2,4]triazin-3(4H)-yl]methyl ester**



- Compound C (660 mg, 2.15 mmol) was dissolved in CH_2Cl_2 (10 mL) and m-chloroperoxybenzoic acid (57%, 745 mg, 2.46 mmol) was added with $MgSO_4$ (2.0 g) and the reaction mixture was stirred at rt for 5 hr. The mixture was filtered and the filtrate was washed with aqueous $NaHCO_3$ solution twice, dried ($MgSO_4$), and concentrated to provide 680 mg (98%) of compound D as a solid. MS: $[M+H]^+ = 324$.

20 **E) 2,2-Dimethylpropanoic acid [5-methoxy-4-oxo-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazin-3(4H)-yl]methyl ester**

- Compound D (680 mg, 2.10 mmol, 1 eq) was dissolved in acetone (10 mL) followed by the addition of benzyl bromide (430 mg, 2.5 mmol) and K_2CO_3 (1.0 g, 7.25 mmol). The reaction mixture was stirred at 60°C for 10 hr, cooled to rt, and filtered. The filtrate was concentrated and purified by flash silica gel chromatography eluting with 25% ethyl acetate in hexanes to provide 485 mg (60%) of E as a gel. MS: $[M+H]^+ = 386$;

F) 5-Methoxy-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazin-4(3H)-one

- 30 Compound E (65 mg, 0.17 mmol) was stirred at rt in a mixture of methanol (1 mL) and ammonium hydroxide (0.2 mL) for 6 hrs. The mixture was concentrated *in vacuo*, dissolved in CH_2Cl_2 , and purified by flash silica gel chromatography eluting with

33% ethyl acetate in hexanes to provide 45 mg (97%) of compound F as a solid. MS: $[M+H]^+ = 272$.


G) 4-Chloro-5-methoxy-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazine

- 5 Compound F (44 mg, 0.16 mmol) was stirred with $POCl_3$ (0.5 mL) at 60°C for 3 hr. The mixture was concentrated *in vacuo*, dissolved in CH_2Cl_2 (2 mL), and stirred with solid $NaHCO_3$ for 10 min. The mixture was filtered and concentrated to provide 46 mg (99%) of compound G as a solid. MS: $[M+H]^+ = 286$ (replacement of Cl by OCH_3 upon standing in methanol); R.T.=3.265 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4 mL/min, monitoring at 220 nm); 1H NMR ($CDCl_3$): δ 8.01 (s, 1H), 7.45-7.30 (m, 6 H), 5.15 (s, 2H), 4.03 (s, 3H).

H) 1,3-Dihydro-3-[5-methoxy-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one

- 15 To a suspension of NaH (60% in oil, 19.2 mg, 0.48 mmol) in of N,N-dimethylformamide (0.5 mL) was added oxindole (63.4 mg, 0.48 mmol). The reaction mixture was stirred for 1 hr at rt. Compound G (38 mg, 0.16 mmol, 1 eq) was added, and the mixture was stirred for 1 hr more. The mixture was diluted with water and filtered. The resulting solid material was triturated with methanol and dried to provide 20 38 mg (62%) of the title compound. MS: $[M+H]^+ = 387$; 1H NMR (*d*-DMSO): δ 12.83 (br s, 1H), 10.64 (br s, 1H), 7.78 (s, 1H), 7.60 (s, 1H), 7.50-7.31 (m, 6H), 7.02-6.94 (m, 1H), 6.89-6.82 (m, 2H), 5.10 (s, 2H), 3.55 (s, 3H).

Example 138



- 25 **4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester**

A) 3-Propylpyrrole-2,4-dicarboxylic acid diethyl ester

- Ethyl isocyanoacetate (4.52 g, 40.0 mmol) was combined with 1,8-diazabicyclo[5.4.0]undec-7-ene (6.09 g, 40.0 mmol) in THF (120 mL). The mixture 30 was warmed to 45°C and butyraldehyde (1.44 g, 20.0 mmol) was added in THF (120

mL) over 30 min. The reaction mixture was stirred at 50°C for 1.5 hr and then was allowed to cool to rt overnight. The solvent was removed *in vacuo* and the resulting brown oil was dissolved in of ethyl acetate (100 mL) and washed with of water (75 mL). The aqueous layer was extracted with ethyl acetate (2 x 100 mL). The combined
5 organic extracts were washed with 0.1 N aqueous HCl (2 x 100 mL) and water (75 mL), dried (Na₂SO₄) and concentrated *in vacuo* to provide 4.270 g (84%) of compound A as a brown oil which was used without further purification. ¹H NMR (CDCl₃): δ 9.17 (br s, 1H), 7.47 (s, 1H), 4.33 (q, *J*=7.1 Hz, 2H), 4.27 (q, *J*=7.1 Hz, 2H), 3.07-3.04 (m, 2H), 1.85-1.78 (m, 2H), 1.39-1.26 (m, 6H), 0.98-0.95 (m, 3H).

10 **B) 1-Amino-3-propylpyrrole-2,4-dicarboxylic acid diethyl ester**

To a suspension of NaH (60% in oil, 0.96 g, 24 mmol) in N,N-dimethylformamide (100 mL) was added compound A (3.06 g, 12 mmol). After 20 min at rt, diphenyl phosphoryl hydroxylamine (5.56 g, 24 mmol) was added, and the mixture was stirred for additional 3 hr. The solvent was removed *in vacuo*, and the residue was dissolved
15 in ethyl acetate and washed with pH 7 phosphate buffer. The organic extracts were dried (Na₂SO₄) and purified by flash chromatography on silica gel eluting with 20% ethyl acetate in hexanes to provide 2 g (63%) of compound B. MS: [M+H]⁺ = 269.2.

C) 5-Propylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one-6-carboxylic acid ethyl ester


20 Compound B (2 g, 7 mmol) was combined with formamide (4.53 g, 100 mmol) and stirred at 160°C for 7 hr. The reaction mixture was allowed to cool to rt. Ice was added to the mixture and the resulting precipitate was filtered off and dried to afford 1.6 g (86%) of compound C. MS: [M-H]⁻ = 247.9.

25 **D) 4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester**

Compound C (19 mg, 0.076 mmol) was stirred with POCl₃ (0.5 mL) at 100°C for 4 hr under Argon. The reaction mixture was concentrated *in vacuo*. CH₃CN (2.50 mL) was added followed by 5-amino-1-benzyl-1H-indazole (25 mg, 0.114 mmol). After 16 hr at rt the mixture was diluted with of ethyl acetate (50 mL) and washed with saturated
30 aqueous NaHCO₃. The combined organic washes were dried (Na₂SO₄), concentrated *in vacuo*, and purified by flash chromatography on silica gel eluting with 40% ethyl acetate in hexanes to provide 20 mg (58%) of the title compound as a yellow-brown

oil. MS: $[M+H]^+ = 455.2$; 1H NMR ($CDCl_3$): δ 8.11 (s, 1H), 8.07 (s, 1H), 8.02 (s, 1H), 7.94 (s, 1H), 7.45-7.19 (m, 7H), 5.61 (s, 2H), 4.35 (q, $J=7.2$ Hz, 2H), 3.29 (t, $J=8.2$ Hz, 2H), 1.85-1.78 (m, 2H), 1.39 (t, $J=7.2$ Hz, 3H), 1.11 (t, $J=7.2$ Hz, 3H).

Example 139



4-[(4-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

A) 3-Ethylpyrrole-2,4-dicarboxylic acid diethyl ester

Ethyl isocyanoacetate (50.67 g, 0.448 mol, 2 eq) was combined with 1,8-diazabicyclo[5.4.0]undec-7-ene (68.2 g, 0.45 mol) in 1.0 L of tetrahydrofuran. The mixture was warmed to 50°C and propanal (13 g, 0.224 mol, 1 eq) was added in 250 mL of tetrahydrofuran. The reaction was stirred at 50°C for 2 hrs. The mixture was allowed to cool to rt and stirred overnight. The solvent was removed *in vacuo*. The resulting brown oil began to crystallize upon standing. The material was triturated with ether, and the resulting solid was collected by filtration and dried to provide (14.0 g, 26%) of compound A as a brown solid. Additional material (18.3 g, 34%) was obtained upon concentration of the mother liquor. MS: $[M+H]^+ = 240$.

B) 5-Ethylpyrrolo[2,1-f][1,2,4]triazin-4(3H)-one-6-carboxylic acid ethyl ester


Compound A (13.8 g, 57.8 mmol) was converted to compound B (8 g, 58%) as a pale yellow solid using the procedure described for the preparation of compound C from compound A in Example 138. MS: $[M+H]^+ = 235.0$.

C) 4-[(4-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester

Compound B (210 mg, 0.893 mmol) was stirred with $POCl_3$ (5 mL) at 100°C for 5.5 hr under Argon. The reaction mixture was concentrated *in vacuo*. Toluene (5 mL) was added to the residue and then removed *in vacuo*. CH_3CN (5 mL) was then added followed by 4-bromoaniline (460 mg, 2.68 mmol). After 14 hr at rt the reaction mixture was poured into saturated aqueous $NaHCO_3$ and extracted with ethyl acetate (3 x 75 mL). The combined organic washes were dried (Na_2SO_4), concentrated *in*

vacuo, and purified by flash chromatography on silica gel eluting with 40% ethyl acetate in hexanes to provide 320 mg (92%) of the title compound as a white crystalline solid. MS: $[M+H]^+ = 389.1$; $^1\text{H NMR}$ (CDCl_3): δ 7.99 (s, 1H), 7.94 (s, 1H), 7.58 (d, $J=8.8$ Hz, 1H), 7.49 (d, $J=8.8$ Hz, 1H), 4.35 (q, $J=7.0$ Hz, 2H), 3.29 (t, $J=7.9$ Hz, 2H), 1.41-1.37 (m, 6H).


Example 140



4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester


The title compound was prepared from compound B of Example 139 using the process described for the preparation of example 139 except 5-amino-1-benzyl-1H-indazole was used to add to the chloroimidate. MS: $(M+H)^+ = 441.24$

Example 141



5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester

Sodium methoxide was generated by addition of NaH (0.163 g, 6.8 mmol) to anhydrous methanol (15 mL) under Argon at 0°C. The mixture was stirred at 0°C for 20 min. Example 140 (30 mg, 0.068 mmol) was added in one portion and the resulting mixture was stirred at rt for 18 hr. The reaction mixture was poured into 50 mL pH 7 phosphate buffer. The aqueous phase was extracted with ethyl acetate (2 x 75 mL). The combined organic washes were dried (Na_2SO_4), concentrated *in vacuo*, and purified by flash chromatography on silica gel eluting with 40% ethyl acetate in hexanes to provide 17.8 mg (61%) of the title compound as a white solid. MS: $[M+H]^+ = 427.2$; $^1\text{H NMR}$ (CDCl_3): 8.09 (s, 1H), 8.06 (s, 1H), 8.00 (s, 1H), 7.94 (s, 1H), 7.46-7.19 (m, 7H), 5.61 (s, 2H), 3.89 (s, 3H), 3.33 (q, $J=7.7$ Hz, 2H), 1.42 (t, $J=7.7$ Hz, 6H).

Example 142

N,5-Diethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

5 A. 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid

Example 140 (320 mg, 0.726 mmol) was dissolved in a mixture of THF (6 mL), methanol (2 mL), and water (2 mL). LiOH (305 mg, 7.26 mmol) was added and the reaction mixture was stirred at 50°C for 24 hr. The mixture was poured into 125 mL pH 4 phosphate buffer and extracted with ethyl acetate (3 x 125 mL). The combined organic washes were dried (Na₂SO₄) and concentrated *in vacuo* to provide a quantitative yield of compound A as a white solid. MS: [M+H]⁺ = 413.

B. N,5-Diethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide

Compound A (31 mg, 0.075 mmol) was dissolved under Argon in a mixture of N,N-dimethylformamide (1.5 mL) and CH₃CN (1.5 mL). Ethylamine (2.0 M in THF, 38 µL, 0.075 mmol) and 1-[3-dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (14 mg, 0.075 mmol) were added, and the reaction was stirred at rt for 18 hr. The mixture was poured into 50 mL of water and extracted with ethyl acetate (2 x 75 mL). The combined organic washes were dried (Na₂SO₄), concentrated *in vacuo*, and purified by flash chromatography on silica gel eluting with 75% ethyl acetate in hexanes to provide 11 mg (33%) of the title compound as a white film. MS: [M+H]⁺ = 440.2; ¹H NMR (CDCl₃): δ 8.10 (s, 1H), 8.06 (s, 1H), 7.93 (s, 1H), 7.71 (s, 1H), 7.46-7.19 (m, 7H), 5.87 (br s, 1H), 5.61 (s, 2H), 3.52-3.45 (m, 2H), 3.30 (q, J=7.5 Hz, 2H), 1.44 (t, J=7.5 Hz, 3H), 1.25 (t, J=7.5 Hz, 3H).

The compounds named in Example 143 were prepared using methods analogous to the procedures described hereinbefore.

Example 143

- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-[[[3-(dimethylamino)propyl]amino]carbonyl]-N-ethyl-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(4-phenoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-4-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 10 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[2-(4-morpholinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-[[[3-(dimethylamino)propyl]amino]carbonyl]-N-ethyl-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 15 4-[(3-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(4-phenoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 20 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-4-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(6-Cyano-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 25 4-(2,5-Dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 3-(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 3-(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[2-(4-morpholinyl)ethyl]urea;
- 30 N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[3-(4-morpholinyl)propyl]urea;

- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[4-(4-morpholinyl)butyl]urea;
- 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-methanol;
- 5 5-Ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-N-[2-(4-morpholinyl)ethyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 10 5-Ethyl-N-[6-[3-(1H-imidazol-1-yl)propyl]-2-pyridinyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 7-Bromo-5-ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 15 7-Bromo-5-ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-methanol;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-4-morpholinebutanamide;
- 5-Ethyl-6-[[2-(4-morpholinyl)ethoxy]methyl]-N-[1-(phenylmethyl)-1H-indazol-5-yl]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
- 20 7-Bromo-5-ethyl-N-[2-(4-morpholinyl)ethyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-2-methylpropanamide;
- 25 3-[6-(Dimethylamino)-7-(hydroxymethyl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl]-1,3-dihydro-2H-indol-2-one;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]methanesulfonamide;
- 3-(5,6-Dimethoxypyrrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 30 N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-4-morpholinepropanesulfonamide;
- [4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxypyrrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;

- 1,3-Dihydro-3-[5-methoxy-6-[[4-(4-methyl-1-piperazinyl)butyl]amino]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 3-(6-Amino-5-methoxypyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 1,3-Dihydro-3-[5-methoxy-6-[[4-(4-morpholinyl)butyl]amino]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 5 4-[(3-Hydroxy-5-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(4-Ethyl-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 5-[(5,6-Dimethoxypyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
- 4-[(4-Bromo-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-Hydroxy-4-(1-methylethyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 5-[(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 20 4-[(4-Hydroxy-2-naphthalenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(4-Carboxy-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Chloro-4-fluorophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
- 25 carboxylic acid methyl ester;
- 4-[(3-Chloro-4-fluorophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 1,3-Dihydro-3-(6-methoxy-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-2H-indol-2-one;
- 4-[(3-Chloro-4-fluorophenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid
- 30 methyl ester;
- 4-[3-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-1-oxopropyl]morpholine;

- 1-[3-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-1-oxopropyl]-4-methylpiperazine;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5,6-dimethoxypyrrolo[2,1-f][1,2,4]triazine-7-carboxylic acid methyl ester;
- 5 4-[(4-Butyl-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
 4-[(3-Hydroxy-4-propylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 4-[(3-Hydroxy-4-methylphenyl)amino]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-(2-methoxyethyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-(3-methoxypropyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 15 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[(tetrahydro-2-furanyl)methyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
 5-Ethoxy-4-[(3-hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-
- 20 carboxylic acid ethyl ester;
 4-[(4-Ethyl-3-hydroxyphenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-[(4-Bromo-3-hydroxyphenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 5-Ethyl-4-(phenylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 5-Ethyl-4-(methylphenylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 5-Ethyl-4-(1,2,3,4-tetrahydro-2-isoquinoliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 30 4-[(3-Hydroxy-4-methylphenyl)amino]-N-(3-methoxypropyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(4-methyl-1-piperazinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;

- 4-[(3-Hydroxy-4-methylphenyl)amino]-N,5-dimethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- N-[2-(Dimethylamino)ethyl]-4-[(3-hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5 5-Ethyl-4-(1,2,3,4-tetrahydro-1-quinoliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(phenylmethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-ethoxypyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid ethyl ester;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-4-[(2-phenylethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 15 N-[4-(Dimethylamino)butyl]-4-[(3-hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-(1-methylethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-
- 20 (methylsulfonyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Chloro-4-fluorophenyl)methyl]-5-(1-methylethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- [4-[(3-Chloro-4-fluorophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;
- 25 5-(1-Methylethyl)-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(Butylamino)-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester
- 5-Ethyl-4-[(2-methoxyethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 30 5-Ethyl-4-(4-morpholinyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[[3-(1H-imidazol-1-yl)propyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;

- 2-Methyl-5-[[5-methyl-6-[3-(2H-1,2,3-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]amino]phenol;
- 5-Ethyl-4-[[[(1S)-1-phenylethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5 5-Ethyl-4-[[[(1R)-1-phenylethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[[2-(2-pyridinyl)ethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(4-Cyano-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid methyl ester;
- 4-[(Cyclohexylmethyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[[4-Cyanocyclohexyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 15 4-[(3-Chloro-4-fluorophenyl)amino]-5-(phenylmethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-(Phenylmethyl)-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-(phenylmethyl)pyrrolo[2,1-f][1,2,4]triazine-
- 20 6-carboxylic acid ethyl ester;
- 4-[[4-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(trans-4-hydroxycyclohexyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 25 N-(3-Bromophenyl)-5-methyl-6-[3-(2H-1,2,3-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
- N-(4-Bromo-2-fluorophenyl)-5-methyl-6-[3-(2H-1,2,3-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
- 1,3-Dihydro-3-[5-methyl-6-[3-(2H-1,2,4-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 30 5-Ethyl-4-[[[(1-hydroxycyclohexyl)methyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;


- 4-[[[(3-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[[[(2-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5 5-Ethyl-N,N-dimethyl-4-[[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-N-[3-(4-morpholinyl)propyl]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-propyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 10 N-(3-Bromophenyl)-5-methyl-6-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
- 4-[(3-Chloro-4-fluorophenyl)amino]-5-[3-(phenylmethoxy)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester; and
- 15 4-[[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester.

Claims

What is claimed is:

1. A compound of formula I

5



its enantiomers, diastereomers, and pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein

- 10 X and Y are independently selected from O, OCO, S, SO, SO₂, CO, CO₂, NR¹⁰, NR¹¹CO, NR¹²CONR¹³, NR¹⁴CO₂, NR¹⁵SO₂, NR¹⁶SO₂NR¹⁷, SO₂NR¹⁸, CONR¹⁹, halogen, nitro, cyano, or X or Y are absent;
- Z is selected from O, S, N, or CR²⁰;
- R¹ is hydrogen, CH₃, OH, OCH₃, SH, SCH₃, OCOR²¹, SOR²², SO₂R²³, SO₂NR²⁴R²⁵,
 15 CO₂R²⁶, CONR²⁷R²⁸, NH₂, NR²⁹SO₂NR³⁰R³¹, NR³²SO₂R³³, NR³⁴COR³⁵, NR³⁶CO₂R³⁷, NR³⁸CONR³⁹R⁴⁰, halogen, nitro, or cyano;
- R² and R³ are independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heterocyclo, substituted heterocyclo, aralkyl, substituted aralkyl, heterocycloalkyl or
 20 substituted heterocycloalkyl, or when X is halo, nitro or cyano R² is absent or when Y is halo, nitro or cyano R³ is absent;
- R⁴ and R⁵ are independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclo, substituted heterocyclo, aralkyl, substituted aralkyl, or R⁴ and R⁵ may together form an optionally substituted monocyclic 5-7 membered
 25 saturated or unsaturated carbocyclic or heterocyclic ring, or an optionally substituted bicyclic 7-11 membered saturated or unsaturated carbocyclic or heterocyclic ring, except that when Z is O or S, R⁵ is absent, or when Z is nitrogen, R⁴ and R⁵ are not both hydrogen;
- R⁶ is hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclo, substituted

- heterocyclo, NR^7R^8 , OR^9 or halogen;
 $\text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{21}, \text{R}^{24}, \text{R}^{25}, \text{R}^{26}, \text{R}^{27}, \text{R}^{28},$
 $\text{R}^{29}, \text{R}^{30}, \text{R}^{31}, \text{R}^{32}, \text{R}^{34}, \text{R}^{35}, \text{R}^{36}, \text{R}^{38}, \text{R}^{39}$ and R^{40} are independently selected
from the group consisting of hydrogen, alkyl, substituted alkyl, aryl,
5 substituted aryl, heterocyclo, or substituted heterocyclo;
 $\text{R}^{22}, \text{R}^{23}, \text{R}^{33}$ and R^{37} are independently selected from the group consisting of alkyl,
substituted alkyl, aryl, substituted aryl, heterocyclo, or substituted
heterocyclo; and
 R^{20} is selected from the group consisting of hydrogen, lower alkyl or substituted
10 alkyl, or R^{20} is absent if the carbon to which it is attached is part of an
unsaturated aryl or heteroaryl ring;
with the provisos that:
a. R^2 may not be hydrogen if X is SO , SO_2 , $\text{NR}^{13}\text{CO}_2$, or $\text{NR}^{14}\text{SO}_2$
b. R^3 may not be hydrogen if Y is SO , SO_2 , $\text{NR}^{13}\text{CO}_2$, or $\text{NR}^{14}\text{SO}_2$.
15
2. The compound according to Claim 1, selected from the group
consisting of:
N-(4-Chlorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine;
2-Methyl-5-(pyrrolo[2,1-f][1,2,4]triazin-4-ylamino)phenol;
20 7-Bromo-N-(4-chlorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine;
N-(4-Chloro-2-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine;
2-Methyl-5-[(6-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol;
N-(4-Bromo-2-fluorophenyl)pyrrolo[2,1-f][1,2,4]triazin-4-amine;
2-Methyl-5-[(5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol;
25 5-[[2-(Dimethylamino)pyrrolo[2,1-f][1,2,4]triazin-4-yl]amino]-2-methylphenol;
5-[(5,7-Dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
5-[(6-Ethyl-5,7-dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
5-[(5,6-Dimethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
2-Methyl-5-[(7-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]phenol;
30 N-(4-Bromo-2-fluorophenyl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine;
N-(4-Bromo-2-fluorophenyl)-7-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine;
1-[2,3-Dihydro-6-(pyrrolo[2,1-f][1,2,4]triazin-4-ylamino)-1H-indol-1-yl]ethanone;
N-(4-Bromo-2-fluorophenyl)-6-methylpyrrolo[2,1-f][1,2,4]triazin-4-amine;

- 5-[(5-Ethylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 carboxylic acid methyl ester;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 5 methanol;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 carboxylic acid;
 1-[[4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-
 yl]carbonyl]-4-methylpiperazine;
 10 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[2-(1-
 pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-phenylpyrrolo[2,1-f][1,2,4]triazine-6-
 carboxylic acid ethyl ester;
 4-[(3-Methoxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 15 carboxylic acid methyl ester;
 2-Methyl-5-[5-methyl-6-(1H-1,2,4-triazol-1-ylmethyl)pyrrolo[2,1-f][1,2,4]triazin-4-
 yl]amino]phenol;
 4-[(3-Cyanophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid
 methyl ester;
 20 4-[(3-Carboxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid
 methyl ester;
 4-[3-(Acetylamino)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic
 acid methyl ester;
 4-[(3-Fluoro-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 25 carboxylic acid methyl ester;
 5-Methyl-4-[(4-methyl-3-nitrophenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-
 carboxylic acid methyl ester;
 5-Methyl-4-[[4-methyl-3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-
 f][1,2,4]triazine-6-carboxylic acid methyl ester;
 30 4-[(3-Cyano-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
 carboxylic acid methyl ester;
 4-[(1-Acetyl-2,3-dihydro-1H-indol-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-
 6-carboxylic acid methyl ester;

- 4-(2,3-Dihydro-1H-indol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Amino-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5 4-[(4-Bromo-2-fluorophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(2,3-Dihydro-1H-indol-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Methyl-4-(5-quinolinylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3,4-Dimethylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 4-[[5-(Hydroxymethyl)-2-methylphenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(1H-Indazol-6-ylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(4-Bromo-3-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 20 4-[(4-Chloro-3-nitrophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(2,3-Dihydro-1,4-benzodioxin-6-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 25 5-Methyl-4-[(5-methyl-2-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(1H-Benzotriazol-5-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Methyl-4-[[2-methyl-5-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 30 4-[(1,2-Dihydro-4-methyl-2-oxo-7-quinoliny)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;

- 4-[(3-Hydroxy-4-methylphenyl)methylamino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5 4-[[2-(Aminosulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- [4-[(3-Hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid methyl ester;
- 4-[(3-Hydroxy-4-methoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Hydroxy-2-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 4-(2-Benzothiazolylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-Hydroxy-4-[(phenylsulfonyl)amino]phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-(Aminosulfonyl)-4-methylphenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-
- 20 6-carboxylic acid methyl ester;
- 4-[(2,3-Dihydro-1,3-dioxo-1H-isoindol-5-yl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-(Hydroxymethyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 25 4-(1H-Indol-6-ylamino)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-[(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
- 4-[(3-Carboxy-4-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 30 4-(3-Hydroxyphenoxy)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Methyl-4-[(4-phenoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;

- 5-Methyl-4-[(1,2,3,4-tetrahydro-1,4-dioxo-6-phthalazinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-(Hydroxysulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5 4-[[3-[(Butylamino)sulfonyl]phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[3-(Aminosulfonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid ethyl ester;
- 4-[[3-(Hydrazinocarbonyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Methyl-4-[[3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 5-Ethyl-4-[(3-hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-
- 20 carboxylic acid ethyl ester;
- 4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 4-[(4-Bromo-2-fluorophenyl)amino]-5-methoxy-N-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(6-Methoxy-3-pyridinyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-
- 30 carboxylic acid ethyl ester;
- 5-Methoxy-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;

- 5-Ethyl-4-[(4-methoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(3-Amino-4-methylphenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5 4-[(3-Hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-5,6-dicarboxylic acid diethyl ester;
- 5-Ethyl-4-[[4-methyl-3-[(methylsulfonyl)amino]phenyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid;
- 10 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-benzimidazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 5-Methyl-4-(1,2,3,4-tetrahydro-3-oxo-1-quinoxaliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Methyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[[6-(Acetyl-amino)-3-pyridinyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 20 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 4-[(6-Chloro-3-pyridinyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-(1H-indazol-6-ylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(6-Methoxy-3-pyridinyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 30 [4-[(6-Methoxy-3-pyridinyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;

- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid;
- 4-Hydroxy-5-methoxypyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[2,3-Dihydro-5-[(methylsulfonyl)amino]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Ethyl-4-[(4-hydroxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 5-Methyl-4-(1H-pyrazol-3-ylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(4-Methoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3,4-Dimethoxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 5-Methyl-4-[(5-methyl-1H-pyrazol-3-yl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-3-oxo-1H-indazol-2-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 20 4-(2,3-Dihydro-3-oxo-1H-indazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[6-Fluoro-2-hydroxy-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[6-Bromo-2-hydroxy-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 25 4-[5-(Aminosulfonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[2,3-Dihydro-5-[[[2-(4-morpholinyl)ethyl]amino]sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 30 4-[2,3-Dihydro-2-oxo-5-[[[2-(1-pyrrolidinyl)ethyl]amino]sulfonyl]-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(5-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;

- 4-[2,3-Dihydro-2-oxo-6-(trifluoromethyl)-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)pyrrolo[2,1-f][1,2,4]triazine-5,6-dicarboxylic acid diethyl ester;
- 5 4-[2,3-Dihydro-6-(methylsulfonyl)-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[5-[(Dimethylamino)sulfonyl]-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[5-(Aminocarbonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 4-[(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;
- 4-(5-Cyano-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 15 4-(2,3-Dihydro-6-methyl-2-oxo-1H-pyrazolo[2,3-d]pyrimidin-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-pyrazolo[2,3-b]pyridin-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 20 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[2-(4-morpholinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-[[[3-(dimethylamino)propyl]amino]carbonyl]-N-ethyl-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 30 4-[2,3-Dihydro-5-[[2-(hydroxyethyl)amino]sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[2,3-Dihydro-5-(4-morpholinylsulfonyl)-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;

- 4-[2,3-Dihydro-5-[(methylamino)sulfonyl]-2-oxo-1H-indol-3-yl]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5 5-Ethyl-4-[(4-phenoxyphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-4-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(6-Cyano-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 4-(2,5-Dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 3-(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 3-(6-Amino-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 15 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[2-(4-morpholinyl)ethyl]urea;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[3-(4-morpholinyl)propyl]urea;
- 20 N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-N'-[4-(4-morpholinyl)butyl]urea;
- 5-Ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-methanol;
- 25 4-(2,3-Dihydro-1-methyl-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-4-[(6-methoxy-3-pyridinyl)amino]-N-methyl-N-[2-(1-pyrrolidinyl)ethyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 30 5-Ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;

- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-methyl-1-piperazinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1H-1,2,3-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(2H-1,2,3-triazol-2-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(1H-1,2,4-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 10 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(2-methyl-1H-imidazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[4-(4-morpholinyl)butyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 15 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(6-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethyl-N-[2-(4-morpholinyl)ethyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 20 5-Ethyl-N-[6-[3-(1H-imidazol-1-yl)propyl]-2-pyridinyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(5-Fluoro-2,3-dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 4-(3-Hydroxy-4-methoxyphenyl)-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 7-Bromo-5-ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 7-Bromo-5-ethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-methanol;
- 30 N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-4-morpholinebutanamide;

- 5-Ethyl-6-[[2-(4-morpholinyl)ethoxy)methyl]-N-[1-(phenylmethyl)-1H-indazol-5-yl]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
- 7-Bromo-5-ethyl-N-[2-(4-morpholinyl)ethyl]-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N,5-dimethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-N-[3-(1H-1,2,4-triazol-1-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 10 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxy-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(4-Bromophenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 15 N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-2-methylpropanamide;
- 3-[6-(Dimethylamino)-7-(hydroxymethyl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl]-1,3-dihydro-2H-indol-2-one;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]methanesulfonamide;
- 20 3-(5,6-Dimethoxypyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- N-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-4-morpholinepropanesulfonamide;
- [4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methoxypyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;
- 25 4-[5-(Aminosulfonyl)-2,3-dihydro-2-oxo-1H-indol-3-yl]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 1,3-Dihydro-3-[5-methoxy-6-[[4-(4-methyl-1-piperazinyl)butyl]amino]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 30 3-(6-Amino-5-methoxypyrrolo[2,1-f][1,2,4]triazin-4-yl)-1,3-dihydro-2H-indol-2-one;
- 1,3-Dihydro-3-[5-methoxy-6-[[4-(4-morpholinyl)butyl]amino]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;

- 4-[(3-Hydroxy-5-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(4-Ethyl-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 5 1,3-Dihydro-3-[5-methyl-6-[2-(1H-1,2,4-triazol-1-yl)ethoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 5-[(5,6-Dimethoxypyrrolo[2,1-f][1,2,4]triazin-4-yl)amino]-2-methylphenol;
- 4-[(4-Bromo-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 10 4-[[3-Hydroxy-4-(1-methylethyl)phenyl]amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-propanoic acid methyl ester;
- 4-[(4-Hydroxy-2-naphthalenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-
- 15 carboxylic acid methyl ester;
- 4-[(4-Carboxy-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Chloro-4-fluorophenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 20 4-[(3-Chloro-4-fluorophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 1,3-Dihydro-3-(6-methoxy-5-methylpyrrolo[2,1-f][1,2,4]triazin-4-yl)-2H-indol-2-one;
- 4-[(3-Chloro-4-fluorophenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 25 4-[3-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-1-oxopropyl]morpholine;
- 1-[3-[4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yl]-1-oxopropyl]-4-methylpiperazine;
- 4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-
- 30 carboxylic acid ethyl ester;
- 4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;

- 4-[(4-Bromophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5,6-dimethoxypyrrolo[2,1-f][1,2,4]triazine-7-carboxylic acid methyl ester;
- 5 4-[(4-Butyl-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Hydroxy-4-propylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-propylpyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid ethyl ester;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-(2-methoxyethyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-N-(3-methoxypropyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 15 4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-methyl-N-[(tetrahydro-2-furanyl)methyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5-Ethoxy-4-[(3-hydroxy-4-methylphenyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 4-[(4-Ethyl-3-hydroxyphenyl)amino]-5-methyl-N-[3-(4-
- 20 morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(4-Bromo-3-hydroxyphenyl)amino]-5-methyl-N-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(2H-1,2,3-triazol-2-yl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 25 5-Ethyl-4-(phenylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-(methylphenylamino)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 5-Ethyl-4-(1,2,3,4-tetrahydro-2-isoquinoliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 30 4-[(3-Hydroxy-4-methylphenyl)amino]-N-(3-methoxypropyl)-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(4-methyl-1-piperazinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;

- 4-[(3-Hydroxy-4-methylphenyl)amino]-N,5-dimethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
N-[2-(Dimethylamino)ethyl]-4-[(3-hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
- 5 5-Ethyl-4-(1,2,3,4-tetrahydro-1-quinoliny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
5-Ethyl-4-[(phenylmethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
4-(2,3-Dihydro-2-oxo-1H-indol-3-yl)-5-ethoxypyrrolo[2,1-f][1,2,4]triazine-6-
- 10 carboxylic acid ethyl ester;
4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-(1-pyrrolidiny)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
5-Ethyl-4-[(2-phenylethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 15 N-[4-(Dimethylamino)butyl]-4-[(3-hydroxy-4-methylphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
4-[(3-Hydroxy-4-methylphenyl)amino]-5-(1-methylethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
4-[(3-Hydroxy-4-methylphenyl)amino]-5-methyl-N-[3-
- 20 (methylsulfonyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
4-[(3-Chloro-4-fluorophenyl)methyl]-5-(1-methylethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
[4-[(3-Chloro-4-fluorophenyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazin-6-yl]carbamic acid phenylmethyl ester;
- 25 5-(1-Methylethyl)-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
4-(Butylamino)-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester
5-Ethyl-4-[(2-methoxyethyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 30 5-Ethyl-4-(4-morpholiny)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
5-Ethyl-4-[[3-(1H-imidazol-1-yl)propyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;

- 2-Methyl-5-[[5-methyl-6-[3-(2H-1,2,3-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]amino]phenol;
 1,3-Dihydro-3-[5-methoxy-6-(phenylmethoxy)pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;
- 5 5-Ethyl-4-[[[(1S)-1-phenylethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 5-Ethyl-4-[[[(1R)-1-phenylethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 5-Ethyl-4-[[2-(2-pyridinyl)ethyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid
- 10 ethyl ester;
 4-[(4-Cyano-3-hydroxyphenyl)amino]-5-methylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester;
 4-[(Cyclohexylmethyl)amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 15 4-[[[(4-Cyanocyclohexyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 4-[(3-Chloro-4-fluorophenyl)amino]-5-(phenylmethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 5-(Phenylmethyl)-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-
- 20 f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 4-[(3-Hydroxy-4-methylphenyl)amino]-5-(phenylmethyl)pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 4-[[[(4-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
- 25 5-Ethyl-4-[(trans-4-hydroxycyclohexyl)amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
 N-(3-Bromophenyl)-5-methyl-6-[3-(2H-1,2,3-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
 N-(4-Bromo-2-fluorophenyl)-5-methyl-6-[3-(2H-1,2,3-triazol-2-
- 30 yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
 1,3-Dihydro-3-[5-methyl-6-[3-(2H-1,2,4-triazol-2-yl)propoxy]pyrrolo[2,1-f][1,2,4]triazin-4-yl]-2H-indol-2-one;

- 5-Ethyl-4-[[[(1-hydroxycyclohexyl)methyl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
4-[[[(3-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
5 4-[[[(2-Bromophenyl)methyl]amino]-5-ethylpyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid ethyl ester;
5-Ethyl-N,N-dimethyl-4-[[1-(phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
4-[(3-Hydroxy-4-methylphenyl)amino]-N-[3-(4-morpholinyl)propyl]-5-
10 propylpyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
4-[(3-Hydroxy-4-methylphenyl)amino]-5-propyl-N-[3-(1-pyrrolidinyl)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxamide;
N-(3-Bromophenyl)-5-methyl-6-[3-(4-morpholinyl)propyl]pyrrolo[2,1-f][1,2,4]triazin-4-amine;
15 4-[(3-Chloro-4-fluorophenyl)amino]-5-[3-(phenylmethoxy)propyl]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester; and
4-[[1-(Phenylmethyl)-1H-indazol-5-yl]amino]pyrrolo[2,1-f][1,2,4]triazine-6-carboxylic acid methyl ester; and
pharmaceutically acceptable salts thereof.

20

3. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

4. A pharmaceutical composition comprising a compound of Claim 1 in
25 combination with pharmaceutically acceptable carrier and an anti-cancer or cytotoxic agent formulated as a fixed dose.

5. The pharmaceutical composition as recited in claim 4, wherein said anti-cancer or cytotoxic agent is selected from the group consisting of: tamoxifen,
30 toremifen, raloxifene, droloxifene, idoxifene, megestrol acetate, anastrozole, letrozole, borazole, exemestane, flutamide, nilutamide, bicalutamide, cyproterone acetate, goserelin acetate, luproside, finasteride, herceptin, methotrexate,

5-fluorouracil, cytosine arabinoside, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin, mithramycin, cisplatin, carboplatin, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepan, vincristine, taxol, taxotere, etoposide, teniposide, amsacrine, irinotecan, topotecan and
5 an epothilone.

6. A method for producing an antiangiogenic effect which comprises administering to a warm-blooded species in need thereof, an effective antiangiogenic producing amount of a compound of Claim 1.

10

7. A method for producing a vascular permeability reducing effect which comprises administering to a warm-blooded species in need thereof an effective vascular permeability reducing amount of a compound of Claim 1.

15 8. A method of inhibiting protein kinase activity of growth factor receptors which comprises administering to a warm-blooded species in need thereof, an effective protein kinase inhibiting amount of a compound of Claim 1.

20 9. A method of inhibiting tyrosine kinase activity of growth factor receptors which comprises administering to a warm-blooded species in need thereof, an effective tyrosine kinase inhibiting amount of a compound of Claim 1.

25 10. A method for treating proliferative diseases, comprising administering to a warm-blooded species in need thereof, a therapeutically effective amount of a composition of Claim 3.

11. A method for treating cancer, comprising administering to a warm-blooded species in need thereof, a therapeutically effective amount of a composition of Claim 3.

30

12. A method for treating inflammation, comprising administering to a warm-blooded species in need thereof, a therapeutically effective amount of a composition of Claim 3.

13. A method for treating autoimmune diseases, comprising administering to a mammalian species in need thereof, a therapeutically effective amount of a composition of Claim 3.

5

14. A method for treating proliferative diseases, comprising administering to warm-blooded species in need thereof, a therapeutically effective amount of a composition of Claim 4.

10

15. A method for treating cancer, comprising administering to a warm-blooded species in need thereof, a therapeutically effective amount of a composition of Claim 4.

15

16. A method for treating inflammation, comprising administering to a mammalian species in need thereof, a therapeutically effective amount of a composition of Claim 4.

20

17. A method for treating autoimmune diseases, comprising administering to a mammalian species in need thereof, a therapeutically effective amount of a composition of Claim 4.

25

18. A method for treating diseases associated with signal transduction pathways operating through growth factor receptors, which comprises administering to a warm-blooded species in need thereof a therapeutically effective amount of a compound of Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/13420

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/53; C07D 487/04

US CL : 514/243; 544/183

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/243; 544/183

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
STN/CAS, structure search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | Chem. abstr., Vol. 124, No. 21, 20 May 1996 (Columbus, OH, USA), page 1411, column 2, the abstract No. 290158a, OTTER et al., 'Conformational Properties of Purine-like C-Nucleosides.' Nucleosides Nucleotides. 1996, 15(1-3), 793-807 (Eng). | 1 |
| X | Chem. abstr., Vol. 124, No. 21, 20 May 1996 (Columbus, OH, USA), page 1337, column 1, the abstract No. 289464r, QUINTELA et al. 'A Ready One-Pot Preparation for Pyrrolo[2,1-f]-[1,2,4]Triazine and Pyrrolo[5,1-c]Pyrimido[4,5-e][1,2,4]Triazine Derivatives.' Tetrahedron. 1996, 52(8), 3037-3048 (Eng). | 1 |
| X | Chem. abstr., Vol. 122, No. 1, 02 January 1995 (Columbus, OH, USA), page 1113, column 1, the abstract No. 9999c, PATIL et al. 'Synthesis of Pyrrolo[2,1-f][1,2,4]Triazine Congeners of Nucleic Acid Purines via the N-Amination of 2-Substituted Pyrroles.' J. Heterocycl. Chem. 1994, 31(4), 781-186 (Eng). | 1 |
| X | Chem. abstr., Vol. 88, No. 17, 24 April 1978 (Columbus, OH, USA), page 541, column 2, the abstract No. 121113q, NEUNHOEFFER et al. 'Cycloaddition Reactions with Azabenzene, XI. Cycloaddition Reactions with Methoxy- and Dialkylamino-1,2,4-Triazines.' Justus Liebigs Ann. Chem. 1977, (9), 1413-1420 (Ger). | 1 |

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 July 2000 (14.07.2000)

Date of mailing of the international search report

14 AUG 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Richard L. Raymond

Telephone No. (703) 308-1235